

**AP Calculus AB**  
**Grading Rubric**  
**Derivatives, Max, Min, and**  
**Inflection**

**2016 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS**

| $x$ | $f(x)$ | $f'(x)$ | $g(x)$ | $g'(x)$ |
|-----|--------|---------|--------|---------|
| 1   | -6     | 3       | 2      | 8       |
| 2   | 2      | -2      | -3     | 0       |
| 3   | 8      | 7       | 6      | 2       |
| 6   | 4      | 5       | 3      | -1      |

6. The functions  $f$  and  $g$  have continuous second derivatives. The table above gives values of the functions and their derivatives at selected values of  $x$ .

(a) Let  $k(x) = f(g(x))$ . Write an equation for the line tangent to the graph of  $k$  at  $x = 3$ .

(b) Let  $h(x) = \frac{g(x)}{f(x)}$ . Find  $h'(1)$ .

(c) Evaluate  $\int_1^3 f''(2x) dx$ .

---


**STOP**

**END OF EXAM**

**AP<sup>®</sup> CALCULUS AB**  
**2015 SCORING GUIDELINES**

**Question 5**

The figure above shows the graph of  $f'$ , the derivative of a twice-differentiable function  $f$ , on the interval  $[-3, 4]$ . The graph of  $f'$  has horizontal tangents at  $x = -1$ ,  $x = 1$ , and  $x = 3$ . The areas of the regions bounded by the  $x$ -axis and the graph of  $f'$  on the intervals  $[-2, 1]$  and  $[1, 4]$  are 9 and 12, respectively.



- (a) Find all  $x$ -coordinates at which  $f$  has a relative maximum. Give a reason for your answer.
- (b) On what open intervals contained in  $-3 < x < 4$  is the graph of  $f$  both concave down and decreasing? Give a reason for your answer.
- (c) Find the  $x$ -coordinates of all points of inflection for the graph of  $f$ . Give a reason for your answer.
- (d) Given that  $f(1) = 3$ , write an expression for  $f(x)$  that involves an integral. Find  $f(4)$  and  $f(-2)$ .

- (a)  $f'(x) = 0$  at  $x = -2$ ,  $x = 1$ , and  $x = 4$ .  
 $f'(x)$  changes from positive to negative at  $x = -2$ .  
 Therefore,  $f$  has a relative maximum at  $x = -2$ .
- (b) The graph of  $f$  is concave down and decreasing on the intervals  $-2 < x < -1$  and  $1 < x < 3$  because  $f'$  is decreasing and negative on these intervals.
- (c) The graph of  $f$  has a point of inflection at  $x = -1$  and  $x = 3$  because  $f'$  changes from decreasing to increasing at these points.

The graph of  $f$  has a point of inflection at  $x = 1$  because  $f'$  changes from increasing to decreasing at this point.

(d)  $f(x) = 3 + \int_1^x f'(t) dt$

$$f(4) = 3 + \int_1^4 f'(t) dt = 3 + (-12) = -9$$

$$\begin{aligned} f(-2) &= 3 + \int_1^{-2} f'(t) dt = 3 - \int_{-2}^1 f'(t) dt \\ &= 3 - (-9) = 12 \end{aligned}$$

$$2 : \begin{cases} 1 : \text{identifies } x = -2 \\ 1 : \text{answer with reason} \end{cases}$$

$$2 : \begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$$

$$2 : \begin{cases} 1 : \text{identifies } x = -1, 1, \text{ and } 3 \\ 1 : \text{reason} \end{cases}$$

$$3 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{expression for } f(x) \\ 1 : f(4) \text{ and } f(-2) \end{cases}$$

**AP<sup>®</sup> CALCULUS AB**  
**2015 SCORING GUIDELINES**

**Question 6**

Consider the curve given by the equation  $y^3 - xy = 2$ . It can be shown that  $\frac{dy}{dx} = \frac{y}{3y^2 - x}$ .

- (a) Write an equation for the line tangent to the curve at the point  $(-1, 1)$ .
- (b) Find the coordinates of all points on the curve at which the line tangent to the curve at that point is vertical.
- (c) Evaluate  $\frac{d^2y}{dx^2}$  at the point on the curve where  $x = -1$  and  $y = 1$ .

(a)  $\left. \frac{dy}{dx} \right|_{(x,y)=(-1,1)} = \frac{1}{3(1)^2 - (-1)} = \frac{1}{4}$

An equation for the tangent line is  $y = \frac{1}{4}(x + 1) + 1$ .

(b)  $3y^2 - x = 0 \Rightarrow x = 3y^2$

So,  $y^3 - xy = 2 \Rightarrow y^3 - (3y^2)(y) = 2 \Rightarrow y = -1$

$(-1)^3 - x(-1) = 2 \Rightarrow x = 3$

The tangent line to the curve is vertical at the point  $(3, -1)$ .

(c)  $\frac{d^2y}{dx^2} = \frac{(3y^2 - x)\frac{dy}{dx} - y\left(6y\frac{dy}{dx} - 1\right)}{(3y^2 - x)^2}$

$\left. \frac{d^2y}{dx^2} \right|_{(x,y)=(-1,1)} = \frac{(3 \cdot 1^2 - (-1)) \cdot \frac{1}{4} - 1 \cdot \left(6 \cdot 1 \cdot \frac{1}{4} - 1\right)}{(3 \cdot 1^2 - (-1))^2}$

$= \frac{1 - \frac{1}{2}}{16} = \frac{1}{32}$

2 :  $\begin{cases} 1 : \text{slope} \\ 1 : \text{equation for tangent line} \end{cases}$

3 :  $\begin{cases} 1 : \text{sets } 3y^2 - x = 0 \\ 1 : \text{equation in one variable} \\ 1 : \text{coordinates} \end{cases}$

4 :  $\begin{cases} 2 : \text{implicit differentiation} \\ 1 : \text{substitution for } \frac{dy}{dx} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2014 SCORING GUIDELINES**

**Question 1**

Grass clippings are placed in a bin, where they decompose. For  $0 \leq t \leq 30$ , the amount of grass clippings remaining in the bin is modeled by  $A(t) = 6.687(0.931)^t$ , where  $A(t)$  is measured in pounds and  $t$  is measured in days.

- (a) Find the average rate of change of  $A(t)$  over the interval  $0 \leq t \leq 30$ . Indicate units of measure.
- (b) Find the value of  $A'(15)$ . Using correct units, interpret the meaning of the value in the context of the problem.
- (c) Find the time  $t$  for which the amount of grass clippings in the bin is equal to the average amount of grass clippings in the bin over the interval  $0 \leq t \leq 30$ .
- (d) For  $t > 30$ ,  $L(t)$ , the linear approximation to  $A$  at  $t = 30$ , is a better model for the amount of grass clippings remaining in the bin. Use  $L(t)$  to predict the time at which there will be 0.5 pound of grass clippings remaining in the bin. Show the work that leads to your answer.

(a)  $\frac{A(30) - A(0)}{30 - 0} = -0.197$  (or  $-0.196$ ) lbs/day

1 : answer with units

(b)  $A'(15) = -0.164$  (or  $-0.163$ )

The amount of grass clippings in the bin is decreasing at a rate of 0.164 (or 0.163) lbs/day at time  $t = 15$  days.

2 :  $\begin{cases} 1 : A'(15) \\ 1 : \text{interpretation} \end{cases}$

(c)  $A(t) = \frac{1}{30} \int_0^{30} A(t) dt \Rightarrow t = 12.415$  (or 12.414)

2 :  $\begin{cases} 1 : \frac{1}{30} \int_0^{30} A(t) dt \\ 1 : \text{answer} \end{cases}$

(d)  $L(t) = A(30) + A'(30) \cdot (t - 30)$

$A'(30) = -0.055976$

$A(30) = 0.782928$


$L(t) = 0.5 \Rightarrow t = 35.054$

4 :  $\begin{cases} 2 : \text{expression for } L(t) \\ 1 : L(t) = 0.5 \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2014 SCORING GUIDELINES**

**Question 3**

The function  $f$  is defined on the closed interval  $[-5, 4]$ . The graph of  $f$  consists of three line segments and is shown in the figure above.



Graph of  $f$

Let  $g$  be the function defined by  $g(x) = \int_{-3}^x f(t) dt$ .

- (a) Find  $g(3)$ .
- (b) On what open intervals contained in  $-5 < x < 4$  is the graph of  $g$  both increasing and concave down? Give a reason for your answer.
- (c) The function  $h$  is defined by  $h(x) = \frac{g(x)}{5x}$ . Find  $h'(3)$ .
- (d) The function  $p$  is defined by  $p(x) = f(x^2 - x)$ . Find the slope of the line tangent to the graph of  $p$  at the point where  $x = -1$ .

(a)  $g(3) = \int_{-3}^3 f(t) dt = 6 + 4 - 1 = 9$

1 : answer

(b)  $g'(x) = f(x)$

2 :  $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

The graph of  $g$  is increasing and concave down on the intervals  $-5 < x < -3$  and  $0 < x < 2$  because  $g' = f$  is positive and decreasing on these intervals.

(c)  $h'(x) = \frac{5xg'(x) - g(x)5}{(5x)^2} = \frac{5xg'(x) - 5g(x)}{25x^2}$

3 :  $\begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$

$$h'(3) = \frac{(5)(3)g'(3) - 5g(3)}{25 \cdot 3^2}$$

$$= \frac{15(-2) - 5(9)}{225} = \frac{-75}{225} = -\frac{1}{3}$$

(d)  $p'(x) = f'(x^2 - x)(2x - 1)$

3 :  $\begin{cases} 2 : p'(x) \\ 1 : \text{answer} \end{cases}$

$$p'(-1) = f'(2)(-3) = (-2)(-3) = 6$$

**AP<sup>®</sup> CALCULUS AB**  
**2014 SCORING GUIDELINES**

**Question 5**

|         |    |               |               |              |   |             |               |
|---------|----|---------------|---------------|--------------|---|-------------|---------------|
| $x$     | -2 | $-2 < x < -1$ | -1            | $-1 < x < 1$ | 1 | $1 < x < 3$ | 3             |
| $f(x)$  | 12 | Positive      | 8             | Positive     | 2 | Positive    | 7             |
| $f'(x)$ | -5 | Negative      | 0             | Negative     | 0 | Positive    | $\frac{1}{2}$ |
| $g(x)$  | -1 | Negative      | 0             | Positive     | 3 | Positive    | 1             |
| $g'(x)$ | 2  | Positive      | $\frac{3}{2}$ | Positive     | 0 | Negative    | -2            |

The twice-differentiable functions  $f$  and  $g$  are defined for all real numbers  $x$ . Values of  $f$ ,  $f'$ ,  $g$ , and  $g'$  for various values of  $x$  are given in the table above.

- (a) Find the  $x$ -coordinate of each relative minimum of  $f$  on the interval  $[-2, 3]$ . Justify your answers.
- (b) Explain why there must be a value  $c$ , for  $-1 < c < 1$ , such that  $f''(c) = 0$ .
- (c) The function  $h$  is defined by  $h(x) = \ln(f(x))$ . Find  $h'(3)$ . Show the computations that lead to your answer.
- (d) Evaluate  $\int_{-2}^3 f'(g(x))g'(x) dx$ .

(a)  $x = 1$  is the only critical point at which  $f'$  changes sign from negative to positive. Therefore,  $f$  has a relative minimum at  $x = 1$ .

(b)  $f'$  is differentiable  $\Rightarrow f'$  is continuous on the interval  $-1 \leq x \leq 1$

$$\frac{f'(1) - f'(-1)}{1 - (-1)} = \frac{0 - 0}{2} = 0$$

Therefore, by the Mean Value Theorem, there is at least one value  $c$ ,  $-1 < c < 1$ , such that  $f''(c) = 0$ .

(c)  $h'(x) = \frac{1}{f(x)} \cdot f'(x)$

$$h'(3) = \frac{1}{f(3)} \cdot f'(3) = \frac{1}{7} \cdot \frac{1}{2} = \frac{1}{14}$$

(d)  $\int_{-2}^3 f'(g(x))g'(x) dx = [f(g(x))]_{x=-2}^{x=3}$   
 $= f(g(3)) - f(g(-2))$   
 $= f(1) - f(-1)$   
 $= 2 - 8 = -6$

1 : answer with justification

$$2 : \begin{cases} 1 : f'(1) - f'(-1) = 0 \\ 1 : \text{explanation, using Mean Value Theorem} \end{cases}$$

$$3 : \begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$$

$$3 : \begin{cases} 2 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB**  
**2013 SCORING GUIDELINES**

**Question 1**

On a certain workday, the rate, in tons per hour, at which unprocessed gravel arrives at a gravel processing plant is modeled by  $G(t) = 90 + 45 \cos\left(\frac{t^2}{18}\right)$ , where  $t$  is measured in hours and  $0 \leq t \leq 8$ . At the beginning of the workday ( $t = 0$ ), the plant has 500 tons of unprocessed gravel. During the hours of operation,  $0 \leq t \leq 8$ , the plant processes gravel at a constant rate of 100 tons per hour.

- (a) Find  $G'(5)$ . Using correct units, interpret your answer in the context of the problem.
- (b) Find the total amount of unprocessed gravel that arrives at the plant during the hours of operation on this workday.
- (c) Is the amount of unprocessed gravel at the plant increasing or decreasing at time  $t = 5$  hours? Show the work that leads to your answer.
- (d) What is the maximum amount of unprocessed gravel at the plant during the hours of operation on this workday? Justify your answer.

(a)  $G'(5) = -24.588$  (or  $-24.587$ )

The rate at which gravel is arriving is decreasing by 24.588 (or 24.587) tons per hour per hour at time  $t = 5$  hours.

(b)  $\int_0^8 G(t) dt = 825.551$  tons

(c)  $G(5) = 98.140764 < 100$

At time  $t = 5$ , the rate at which unprocessed gravel is arriving is less than the rate at which it is being processed. Therefore, the amount of unprocessed gravel at the plant is decreasing at time  $t = 5$ .

- (d) The amount of unprocessed gravel at time  $t$  is given by

$$A(t) = 500 + \int_0^t (G(s) - 100) ds.$$

$$A'(t) = G(t) - 100 = 0 \Rightarrow t = 4.923480$$

| $t$     | $A(t)$     |
|---------|------------|
| 0       | 500        |
| 4.92348 | 635.376123 |
| 8       | 525.551089 |

The maximum amount of unprocessed gravel at the plant during this workday is 635.376 tons.

$$2 : \begin{cases} 1 : G'(5) \\ 1 : \text{interpretation with units} \end{cases}$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

$$2 : \begin{cases} 1 : \text{compares } G(5) \text{ to } 100 \\ 1 : \text{conclusion} \end{cases}$$

$$3 : \begin{cases} 1 : \text{considers } A'(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$



**AP<sup>®</sup> CALCULUS AB**  
**2012 SCORING GUIDELINES**

**Question 4**

The function  $f$  is defined by  $f(x) = \sqrt{25 - x^2}$  for  $-5 \leq x \leq 5$ .

- (a) Find  $f'(x)$ .
- (b) Write an equation for the line tangent to the graph of  $f$  at  $x = -3$ .
- (c) Let  $g$  be the function defined by  $g(x) = \begin{cases} f(x) & \text{for } -5 \leq x \leq -3 \\ x + 7 & \text{for } -3 < x \leq 5. \end{cases}$   
Is  $g$  continuous at  $x = -3$ ? Use the definition of continuity to explain your answer.
- (d) Find the value of  $\int_0^5 x\sqrt{25 - x^2} dx$ .

(a)  $f'(x) = \frac{1}{2}(25 - x^2)^{-1/2}(-2x) = \frac{-x}{\sqrt{25 - x^2}}, \quad -5 < x < 5$

2 :  $f'(x)$

(b)  $f'(-3) = \frac{3}{\sqrt{25 - 9}} = \frac{3}{4}$

$f(-3) = \sqrt{25 - 9} = 4$

An equation for the tangent line is  $y = 4 + \frac{3}{4}(x + 3)$ .

2 :  $\begin{cases} 1 : f'(-3) \\ 1 : \text{answer} \end{cases}$

(c)  $\lim_{x \rightarrow -3^-} g(x) = \lim_{x \rightarrow -3^-} f(x) = \lim_{x \rightarrow -3^-} \sqrt{25 - x^2} = 4$

$\lim_{x \rightarrow -3^+} g(x) = \lim_{x \rightarrow -3^+} (x + 7) = 4$

Therefore,  $\lim_{x \rightarrow -3} g(x) = 4$ .

$g(-3) = f(-3) = 4$

So,  $\lim_{x \rightarrow -3} g(x) = g(-3)$ .

Therefore,  $g$  is continuous at  $x = -3$ .

2 :  $\begin{cases} 1 : \text{considers one-sided limits} \\ 1 : \text{answer with explanation} \end{cases}$

(d) Let  $u = 25 - x^2 \Rightarrow du = -2x dx$

$\int_0^5 x\sqrt{25 - x^2} dx = -\frac{1}{2} \int_{25}^0 \sqrt{u} du$

$= \left[ -\frac{1}{2} \cdot \frac{2}{3} u^{3/2} \right]_{u=25}^{u=0}$

$= -\frac{1}{3}(0 - 125) = \frac{125}{3}$

3 :  $\begin{cases} 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2011 SCORING GUIDELINES**

**Question 6**

Let  $f$  be a function defined by  $f(x) = \begin{cases} 1 - 2\sin x & \text{for } x \leq 0 \\ e^{-4x} & \text{for } x > 0. \end{cases}$

- (a) Show that  $f$  is continuous at  $x = 0$ .  
 (b) For  $x \neq 0$ , express  $f'(x)$  as a piecewise-defined function. Find the value of  $x$  for which  $f'(x) = -3$ .  
 (c) Find the average value of  $f$  on the interval  $[-1, 1]$ .

(a)  $\lim_{x \rightarrow 0^-} (1 - 2\sin x) = 1$

$$\lim_{x \rightarrow 0^+} e^{-4x} = 1$$

$$f(0) = 1$$

So,  $\lim_{x \rightarrow 0} f(x) = f(0)$ .

Therefore  $f$  is continuous at  $x = 0$ .

(b)  $f'(x) = \begin{cases} -2\cos x & \text{for } x < 0 \\ -4e^{-4x} & \text{for } x > 0 \end{cases}$

$$-2\cos x \neq -3 \text{ for all values of } x < 0.$$

$$-4e^{-4x} = -3 \text{ when } x = -\frac{1}{4} \ln\left(\frac{3}{4}\right) > 0.$$

Therefore  $f'(x) = -3$  for  $x = -\frac{1}{4} \ln\left(\frac{3}{4}\right)$ .

(c) 
$$\begin{aligned} \int_{-1}^1 f(x) dx &= \int_{-1}^0 f(x) dx + \int_0^1 f(x) dx \\ &= \int_{-1}^0 (1 - 2\sin x) dx + \int_0^1 e^{-4x} dx \\ &= \left[ x + 2\cos x \right]_{x=-1}^{x=0} + \left[ -\frac{1}{4}e^{-4x} \right]_{x=0}^{x=1} \\ &= (3 - 2\cos(-1)) + \left( -\frac{1}{4}e^{-4} + \frac{1}{4} \right) \end{aligned}$$

$$\begin{aligned} \text{Average value} &= \frac{1}{2} \int_{-1}^1 f(x) dx \\ &= \frac{13}{8} - \cos(-1) - \frac{1}{8}e^{-4} \end{aligned}$$

2 : analysis

3 :  $\begin{cases} 2 : f'(x) \\ 1 : \text{value of } x \end{cases}$

4 :  $\begin{cases} 1 : \int_{-1}^0 (1 - 2\sin x) dx \text{ and } \int_0^1 e^{-4x} dx \\ 2 : \text{antiderivatives} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2010 SCORING GUIDELINES**

**Question 6**

Solutions to the differential equation  $\frac{dy}{dx} = xy^3$  also satisfy  $\frac{d^2y}{dx^2} = y^3(1 + 3x^2y^2)$ . Let  $y = f(x)$  be a particular solution to the differential equation  $\frac{dy}{dx} = xy^3$  with  $f(1) = 2$ .

- (a) Write an equation for the line tangent to the graph of  $y = f(x)$  at  $x = 1$ .
- (b) Use the tangent line equation from part (a) to approximate  $f(1.1)$ . Given that  $f(x) > 0$  for  $1 < x < 1.1$ , is the approximation for  $f(1.1)$  greater than or less than  $f(1.1)$ ? Explain your reasoning.
- (c) Find the particular solution  $y = f(x)$  with initial condition  $f(1) = 2$ .

(a)  $f'(1) = \left. \frac{dy}{dx} \right|_{(1,2)} = 8$

An equation of the tangent line is  $y = 2 + 8(x - 1)$ .

$$2 : \begin{cases} 1 : f'(1) \\ 1 : \text{answer} \end{cases}$$

(b)  $f(1.1) \approx 2.8$

Since  $y = f(x) > 0$  on the interval  $1 \leq x < 1.1$ ,

$$\frac{d^2y}{dx^2} = y^3(1 + 3x^2y^2) > 0 \text{ on this interval.}$$

Therefore on the interval  $1 < x < 1.1$ , the line tangent to the graph of  $y = f(x)$  at  $x = 1$  lies below the curve and the approximation 2.8 is less than  $f(1.1)$ .

$$2 : \begin{cases} 1 : \text{approximation} \\ 1 : \text{conclusion with explanation} \end{cases}$$

(c)  $\frac{dy}{dx} = xy^3$

$$\int \frac{1}{y^3} dy = \int x dx$$

$$-\frac{1}{2y^2} = \frac{x^2}{2} + C$$

$$-\frac{1}{2 \cdot 2^2} = \frac{1^2}{2} + C \Rightarrow C = -\frac{5}{8}$$

$$y^2 = \frac{1}{\frac{5}{4} - x^2}$$

$$f(x) = \frac{2}{\sqrt{5 - 4x^2}}, \quad \frac{-\sqrt{5}}{2} < x < \frac{\sqrt{5}}{2}$$

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

**AP<sup>®</sup> CALCULUS AB**  
**2008 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function given by  $f(x) = \frac{\ln x}{x}$  for all  $x > 0$ . The derivative of  $f$  is given by

$$f'(x) = \frac{1 - \ln x}{x^2}.$$

- (a) Write an equation for the line tangent to the graph of  $f$  at  $x = e^2$ .  
 (b) Find the  $x$ -coordinate of the critical point of  $f$ . Determine whether this point is a relative minimum, a relative maximum, or neither for the function  $f$ . Justify your answer.  
 (c) The graph of the function  $f$  has exactly one point of inflection. Find the  $x$ -coordinate of this point.  
 (d) Find  $\lim_{x \rightarrow 0^+} f(x)$ .

(a)  $f(e^2) = \frac{\ln e^2}{e^2} = \frac{2}{e^2}$ ,  $f'(e^2) = \frac{1 - \ln e^2}{(e^2)^2} = -\frac{1}{e^4}$

An equation for the tangent line is  $y = \frac{2}{e^2} - \frac{1}{e^4}(x - e^2)$ .

2 :  $\begin{cases} 1 : f(e^2) \text{ and } f'(e^2) \\ 1 : \text{answer} \end{cases}$

- (b)  $f'(x) = 0$  when  $x = e$ . The function  $f$  has a relative maximum at  $x = e$  because  $f'(x)$  changes from positive to negative at  $x = e$ .

3 :  $\begin{cases} 1 : x = e \\ 1 : \text{relative maximum} \\ 1 : \text{justification} \end{cases}$

(c)  $f''(x) = \frac{-\frac{1}{x}x^2 - (1 - \ln x)2x}{x^4} = \frac{-3 + 2\ln x}{x^3}$  for all  $x > 0$

$f''(x) = 0$  when  $-3 + 2\ln x = 0$

$x = e^{3/2}$

The graph of  $f$  has a point of inflection at  $x = e^{3/2}$  because  $f''(x)$  changes sign at  $x = e^{3/2}$ .

3 :  $\begin{cases} 2 : f''(x) \\ 1 : \text{answer} \end{cases}$

(d)  $\lim_{x \rightarrow 0^+} \frac{\ln x}{x} = -\infty$  or Does Not Exist

1 : answer

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function defined by  $f(x) = k\sqrt{x} - \ln x$  for  $x > 0$ , where  $k$  is a positive constant.

- (a) Find  $f'(x)$  and  $f''(x)$ .
- (b) For what value of the constant  $k$  does  $f$  have a critical point at  $x = 1$ ? For this value of  $k$ , determine whether  $f$  has a relative minimum, relative maximum, or neither at  $x = 1$ . Justify your answer.
- (c) For a certain value of the constant  $k$ , the graph of  $f$  has a point of inflection on the  $x$ -axis. Find this value of  $k$ .

(a)  $f'(x) = \frac{k}{2\sqrt{x}} - \frac{1}{x}$

$$f''(x) = -\frac{1}{4}kx^{-3/2} + x^{-2}$$

$$2 : \begin{cases} 1 : f'(x) \\ 1 : f''(x) \end{cases}$$

(b)  $f'(1) = \frac{1}{2}k - 1 = 0 \Rightarrow k = 2$

When  $k = 2$ ,  $f'(1) = 0$  and  $f''(1) = -\frac{1}{2} + 1 > 0$ .

$f$  has a relative minimum value at  $x = 1$  by the Second Derivative Test.

$$4 : \begin{cases} 1 : \text{sets } f'(1) = 0 \text{ or } f'(x) = 0 \\ 1 : \text{solves for } k \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

(c) At this inflection point,  $f''(x) = 0$  and  $f(x) = 0$ .

$$f''(x) = 0 \Rightarrow \frac{-k}{4x^{3/2}} + \frac{1}{x^2} = 0 \Rightarrow k = \frac{4}{\sqrt{x}}$$

$$f(x) = 0 \Rightarrow k\sqrt{x} - \ln x = 0 \Rightarrow k = \frac{\ln x}{\sqrt{x}}$$

$$3 : \begin{cases} 1 : f''(x) = 0 \text{ or } f(x) = 0 \\ 1 : \text{equation in one variable} \\ 1 : \text{answer} \end{cases}$$

Therefore,  $\frac{4}{\sqrt{x}} = \frac{\ln x}{\sqrt{x}}$   
 $\Rightarrow 4 = \ln x$   
 $\Rightarrow x = e^4$   
 $\Rightarrow k = \frac{4}{e^2}$

**AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES**

**Question 6**

The twice-differentiable function  $f$  is defined for all real numbers and satisfies the following conditions:

$$f(0) = 2, \quad f'(0) = -4, \quad \text{and} \quad f''(0) = 3.$$

- (a) The function  $g$  is given by  $g(x) = e^{ax} + f(x)$  for all real numbers, where  $a$  is a constant. Find  $g'(0)$  and  $g''(0)$  in terms of  $a$ . Show the work that leads to your answers.
- (b) The function  $h$  is given by  $h(x) = \cos(kx)f(x)$  for all real numbers, where  $k$  is a constant. Find  $h'(x)$  and write an equation for the line tangent to the graph of  $h$  at  $x = 0$ .

(a)  $g'(x) = ae^{ax} + f'(x)$   
 $g'(0) = a - 4$

$g''(x) = a^2e^{ax} + f''(x)$   
 $g''(0) = a^2 + 3$

$$4 : \begin{cases} 1 : g'(x) \\ 1 : g'(0) \\ 1 : g''(x) \\ 1 : g''(0) \end{cases}$$

(b)  $h'(x) = f'(x)\cos(kx) - k\sin(kx)f(x)$   
 $h'(0) = f'(0)\cos(0) - k\sin(0)f(0) = f'(0) = -4$   
 $h(0) = \cos(0)f(0) = 2$   
 The equation of the tangent line is  $y = -4x + 2$ .

$$5 : \begin{cases} 2 : h'(x) \\ 3 : \begin{cases} 1 : h'(0) \\ 1 : h(0) \\ 1 : \text{equation of tangent line} \end{cases} \end{cases}$$

**AP Calculus AB  
Grading Rubric  
FTC I and II**


**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2016 SCORING GUIDELINES**

**Question 3**

The figure above shows the graph of the piecewise-linear function  $f$ . For  $-4 \leq x \leq 12$ , the function  $g$  is defined by

$$g(x) = \int_2^x f(t) dt.$$

- (a) Does  $g$  have a relative minimum, a relative maximum, or neither at  $x = 10$ ? Justify your answer.
- (b) Does the graph of  $g$  have a point of inflection at  $x = 4$ ? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of  $g$  on the interval  $-4 \leq x \leq 12$ . Justify your answers.
- (d) For  $-4 \leq x \leq 12$ , find all intervals for which  $g(x) \leq 0$ .



Graph of  $f$

- (a) The function  $g$  has neither a relative minimum nor a relative maximum at  $x = 10$  since  $g'(x) = f(x)$  and  $f(x) \leq 0$  for  $8 \leq x \leq 12$ .
- (b) The graph of  $g$  has a point of inflection at  $x = 4$  since  $g'(x) = f(x)$  is increasing for  $2 \leq x \leq 4$  and decreasing for  $4 \leq x \leq 8$ .
- (c)  $g'(x) = f(x)$  changes sign only at  $x = -2$  and  $x = 6$ .

| $x$ | $g(x)$ |
|-----|--------|
| -4  | -4     |
| -2  | -8     |
| 6   | 8      |
| 12  | -4     |

On the interval  $-4 \leq x \leq 12$ , the absolute minimum value is  $g(-2) = -8$  and the absolute maximum value is  $g(6) = 8$ .

- (d)  $g(x) \leq 0$  for  $-4 \leq x \leq 2$  and  $10 \leq x \leq 12$ .

1 :  $g'(x) = f(x)$  in (a), (b), (c), or (d)

1 : answer with justification

1 : answer with justification

4 : { 1 : considers  $x = -2$  and  $x = 6$   
as candidates  
1 : considers  $x = -4$  and  $x = 12$   
2 : answers with justification


2 : intervals



**AP<sup>®</sup> CALCULUS AB**  
**2013 SCORING GUIDELINES**

**Question 4**

The figure above shows the graph of  $f'$ , the derivative of a twice-differentiable function  $f$ , on the closed interval  $0 \leq x \leq 8$ . The graph of  $f'$  has horizontal tangent lines at  $x = 1$ ,  $x = 3$ , and  $x = 5$ . The areas of the regions between the graph of  $f'$  and the  $x$ -axis are labeled in the figure. The function  $f$  is defined for all real numbers and satisfies  $f(8) = 4$ .



- (a) Find all values of  $x$  on the open interval  $0 < x < 8$  for which the function  $f$  has a local minimum. Justify your answer.
- (b) Determine the absolute minimum value of  $f$  on the closed interval  $0 \leq x \leq 8$ . Justify your answer.
- (c) On what open intervals contained in  $0 < x < 8$  is the graph of  $f$  both concave down and increasing? Explain your reasoning.
- (d) The function  $g$  is defined by  $g(x) = (f(x))^3$ . If  $f(3) = -\frac{5}{2}$ , find the slope of the line tangent to the graph of  $g$  at  $x = 3$ .

(a)  $x = 6$  is the only critical point at which  $f'$  changes sign from negative to positive. Therefore,  $f$  has a local minimum at  $x = 6$ .

(b) From part (a), the absolute minimum occurs either at  $x = 6$  or at an endpoint.

$$\begin{aligned} f(0) &= f(8) + \int_8^0 f'(x) \, dx \\ &= f(8) - \int_0^8 f'(x) \, dx = 4 - 12 = -8 \end{aligned}$$

$$\begin{aligned} f(6) &= f(8) + \int_8^6 f'(x) \, dx \\ &= f(8) - \int_6^8 f'(x) \, dx = 4 - 7 = -3 \end{aligned}$$

$$f(8) = 4$$

The absolute minimum value of  $f$  on the closed interval  $[0, 8]$  is  $-8$ .

(c) The graph of  $f$  is concave down and increasing on  $0 < x < 1$  and  $3 < x < 4$ , because  $f'$  is decreasing and positive on these intervals.

(d)  $g'(x) = 3[f(x)]^2 \cdot f'(x)$

$$g'(3) = 3[f(3)]^2 \cdot f'(3) = 3\left(-\frac{5}{2}\right)^2 \cdot 4 = 75$$

1 : answer with justification

3 :  $\begin{cases} 1 : \text{considers } x = 0 \text{ and } x = 6 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$


2 :  $\begin{cases} 1 : \text{answer} \\ 1 : \text{explanation} \end{cases}$

3 :  $\begin{cases} 2 : g'(x) \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2012 SCORING GUIDELINES**

**Question 3**

Let  $f$  be the continuous function defined on  $[-4, 3]$  whose graph, consisting of three line segments and a semicircle centered at the origin, is given above. Let  $g$  be the function given by  $g(x) = \int_1^x f(t) dt$ .



Graph of  $f$

- (a) Find the values of  $g(2)$  and  $g(-2)$ .
- (b) For each of  $g'(-3)$  and  $g''(-3)$ , find the value or state that it does not exist.
- (c) Find the  $x$ -coordinate of each point at which the graph of  $g$  has a horizontal tangent line. For each of these points, determine whether  $g$  has a relative minimum, relative maximum, or neither a minimum nor a maximum at the point. Justify your answers.
- (d) For  $-4 < x < 3$ , find all values of  $x$  for which the graph of  $g$  has a point of inflection. Explain your reasoning.

(a)  $g(2) = \int_1^2 f(t) dt = -\frac{1}{2}(1)\left(\frac{1}{2}\right) = -\frac{1}{4}$

$$g(-2) = \int_1^{-2} f(t) dt = -\int_{-2}^1 f(t) dt$$

$$= -\left(\frac{3}{2} - \frac{\pi}{2}\right) = \frac{\pi}{2} - \frac{3}{2}$$

(b)  $g'(x) = f(x) \Rightarrow g'(-3) = f(-3) = 2$   
 $g''(x) = f'(x) \Rightarrow g''(-3) = f'(-3) = 1$

- (c) The graph of  $g$  has a horizontal tangent line where  $g'(x) = f(x) = 0$ . This occurs at  $x = -1$  and  $x = 1$ .

$g'(x)$  changes sign from positive to negative at  $x = -1$ .  
Therefore,  $g$  has a relative maximum at  $x = -1$ .

$g'(x)$  does not change sign at  $x = 1$ . Therefore,  $g$  has neither a relative maximum nor a relative minimum at  $x = 1$ .

- (d) The graph of  $g$  has a point of inflection at each of  $x = -2$ ,  $x = 0$ , and  $x = 1$  because  $g''(x) = f'(x)$  changes sign at each of these values.

$$2 : \begin{cases} 1 : g(2) \\ 1 : g(-2) \end{cases}$$


$$2 : \begin{cases} 1 : g'(-3) \\ 1 : g''(-3) \end{cases}$$

$$3 : \begin{cases} 1 : \text{considers } g'(x) = 0 \\ 1 : x = -1 \text{ and } x = 1 \\ 1 : \text{answers with justifications} \end{cases}$$

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{explanation} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB  
2010 SCORING GUIDELINES**

**Question 5**



The function  $g$  is defined and differentiable on the closed interval  $[-7, 5]$  and satisfies  $g(0) = 5$ . The graph of  $y = g'(x)$ , the derivative of  $g$ , consists of a semicircle and three line segments, as shown in the figure above.

- (a) Find  $g(3)$  and  $g(-2)$ .
- (b) Find the  $x$ -coordinate of each point of inflection of the graph of  $y = g(x)$  on the interval  $-7 < x < 5$ . Explain your reasoning.
- (c) The function  $h$  is defined by  $h(x) = g(x) - \frac{1}{2}x^2$ . Find the  $x$ -coordinate of each critical point of  $h$ , where  $-7 < x < 5$ , and classify each critical point as the location of a relative minimum, relative maximum, or neither a minimum nor a maximum. Explain your reasoning.

(a)  $g(3) = 5 + \int_0^3 g'(x) dx = 5 + \frac{\pi \cdot 2^2}{4} + \frac{3}{2} = \frac{13}{2} + \pi$   
 $g(-2) = 5 + \int_0^{-2} g'(x) dx = 5 - \pi$

$$3 : \begin{cases} 1 : \text{uses } g(0) = 5 \\ 1 : g(3) \\ 1 : g(-2) \end{cases}$$

- (b) The graph of  $y = g(x)$  has points of inflection at  $x = 0$ ,  $x = 2$ , and  $x = 3$  because  $g'$  changes from increasing to decreasing at  $x = 0$  and  $x = 3$ , and  $g'$  changes from decreasing to increasing at  $x = 2$ .

$$2 : \begin{cases} 1 : \text{identifies } x = 0, 2, 3 \\ 1 : \text{explanation} \end{cases}$$

- (c)  $h'(x) = g'(x) - x = 0 \Rightarrow g'(x) = x$   
 On the interval  $-2 \leq x \leq 2$ ,  $g'(x) = \sqrt{4 - x^2}$ .  
 On this interval,  $g'(x) = x$  when  $x = \sqrt{2}$ .  
 The only other solution to  $g'(x) = x$  is  $x = 3$ .

$$4 : \begin{cases} 1 : h'(x) \\ 1 : \text{identifies } x = \sqrt{2}, 3 \\ 1 : \text{answer for } \sqrt{2} \text{ with analysis} \\ 1 : \text{answer for 3 with analysis} \end{cases}$$

$$h'(x) = g'(x) - x > 0 \text{ for } 0 \leq x < \sqrt{2}$$

$$h'(x) = g'(x) - x \leq 0 \text{ for } \sqrt{2} < x \leq 5$$

Therefore  $h$  has a relative maximum at  $x = \sqrt{2}$ , and  $h$  has neither a minimum nor a maximum at  $x = 3$ .


**AP<sup>®</sup> CALCULUS AB**  
**2009 SCORING GUIDELINES**

**Question 6**

The derivative of a function  $f$  is defined by

$$f'(x) = \begin{cases} g(x) & \text{for } -4 \leq x \leq 0 \\ 5e^{-x/3} - 3 & \text{for } 0 < x \leq 4 \end{cases}$$

The graph of the continuous function  $f'$ , shown in the figure above, has  $x$ -intercepts at  $x = -2$  and  $x = 3\ln\left(\frac{5}{3}\right)$ . The graph of  $g$  on  $-4 \leq x \leq 0$  is a semicircle, and  $f(0) = 5$ .



Graph of  $f'$

- (a) For  $-4 < x < 4$ , find all values of  $x$  at which the graph of  $f$  has a point of inflection. Justify your answer.
- (b) Find  $f(-4)$  and  $f(4)$ .
- (c) For  $-4 \leq x \leq 4$ , find the value of  $x$  at which  $f$  has an absolute maximum. Justify your answer.

- (a)  $f'$  changes from decreasing to increasing at  $x = -2$  and from increasing to decreasing at  $x = 0$ . Therefore, the graph of  $f$  has points of inflection at  $x = -2$  and  $x = 0$ .

2 :  $\begin{cases} 1 : \text{identifies } x = -2 \text{ or } x = 0 \\ 1 : \text{answer with justification} \end{cases}$

(b)  $f(-4) = 5 + \int_0^{-4} g(x) dx$   
 $= 5 - (8 - 2\pi) = 2\pi - 3$

$$f(4) = 5 + \int_0^4 (5e^{-x/3} - 3) dx$$

$$= 5 + \left. (-15e^{-x/3} - 3x) \right|_{x=0}^{x=4}$$

$$= 8 - 15e^{-4/3}$$

5 :  $\begin{cases} 2 : f(-4) \\ 1 : \text{integral} \\ 1 : \text{value} \\ 3 : f(4) \\ 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{value} \end{cases}$

- (c) Since  $f'(x) > 0$  on the intervals  $-4 < x < -2$  and  $-2 < x < 3\ln\left(\frac{5}{3}\right)$ ,  $f$  is increasing on the interval  $-4 \leq x \leq 3\ln\left(\frac{5}{3}\right)$ .

Since  $f'(x) < 0$  on the interval  $3\ln\left(\frac{5}{3}\right) < x < 4$ ,  $f$  is decreasing on the interval  $3\ln\left(\frac{5}{3}\right) \leq x \leq 4$ .

Therefore,  $f$  has an absolute maximum at  $x = 3\ln\left(\frac{5}{3}\right)$ .

2 :  $\begin{cases} 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$


AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES

Question 3

The graph of the function  $f$  shown above consists of six line segments. Let  $g$  be the function given by

$$g(x) = \int_0^x f(t) dt.$$

- (a) Find  $g(4)$ ,  $g'(4)$ , and  $g''(4)$ .  
 (b) Does  $g$  have a relative minimum, a relative maximum, or neither at  $x = 1$ ? Justify your answer.



Graph of  $f$

- (c) Suppose that  $f$  is defined for all real numbers  $x$  and is periodic with a period of length 5. The graph above shows two periods of  $f$ . Given that  $g(5) = 2$ , find  $g(10)$  and write an equation for the line tangent to the graph of  $g$  at  $x = 108$ .

(a)  $g(4) = \int_0^4 f(t) dt = 3$

$$g'(4) = f(4) = 0$$

$$g''(4) = f'(4) = -2$$

- (b)  $g$  has a relative minimum at  $x = 1$  because  $g' = f$  changes from negative to positive at  $x = 1$ .

- (c)  $g(0) = 0$  and the function values of  $g$  increase by 2 for every increase of 5 in  $x$ .

$$g(10) = 2g(5) = 4$$

$$\begin{aligned} g(108) &= \int_0^{105} f(t) dt + \int_{105}^{108} f(t) dt \\ &= 21g(5) + g(3) = 44 \end{aligned}$$

$$g'(108) = f(108) = f(3) = 2$$

An equation for the line tangent to the graph of  $g$  at  $x = 108$  is  $y - 44 = 2(x - 108)$ .

$$3 : \begin{cases} 1 : g(4) \\ 1 : g'(4) \\ 1 : g''(4) \end{cases}$$

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$$

$$4 : \begin{cases} 1 : g(10) \\ 3 : \begin{cases} 1 : g(108) \\ 1 : g'(108) \\ 1 : \text{equation of tangent line} \end{cases} \end{cases}$$

**AP Calculus AB**  
**Grading Rubric**  
**Position, Velocity, and**  
**Acceleration**

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC**  
**2015 SCORING GUIDELINES**

**Question 3**

|                               |   |     |     |      |     |
|-------------------------------|---|-----|-----|------|-----|
| $t$<br>(minutes)              | 0 | 12  | 20  | 24   | 40  |
| $v(t)$<br>(meters per minute) | 0 | 200 | 240 | -220 | 150 |

Johanna jogs along a straight path. For  $0 \leq t \leq 40$ , Johanna's velocity is given by a differentiable function  $v$ . Selected values of  $v(t)$ , where  $t$  is measured in minutes and  $v(t)$  is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of  $v'(16)$ .
- (b) Using correct units, explain the meaning of the definite integral  $\int_0^{40} |v(t)| dt$  in the context of the problem.

Approximate the value of  $\int_0^{40} |v(t)| dt$  using a right Riemann sum with the four subintervals indicated in the table.

- (c) Bob is riding his bicycle along the same path. For  $0 \leq t \leq 10$ , Bob's velocity is modeled by  $B(t) = t^3 - 6t^2 + 300$ , where  $t$  is measured in minutes and  $B(t)$  is measured in meters per minute.

Find Bob's acceleration at time  $t = 5$ .

- (d) Based on the model  $B$  from part (c), find Bob's average velocity during the interval  $0 \leq t \leq 10$ .

(a)  $v'(16) \approx \frac{240 - 200}{20 - 12} = 5 \text{ meters/min}^2$

1 : approximation

- (b)  $\int_0^{40} |v(t)| dt$  is the total distance Johanna jogs, in meters, over the time interval  $0 \leq t \leq 40$  minutes.

3 :  $\begin{cases} 1 : \text{explanation} \\ 1 : \text{right Riemann sum} \\ 1 : \text{approximation} \end{cases}$

$$\begin{aligned} \int_0^{40} |v(t)| dt &\approx 12 \cdot |v(12)| + 8 \cdot |v(20)| + 4 \cdot |v(24)| + 16 \cdot |v(40)| \\ &= 12 \cdot 200 + 8 \cdot 240 + 4 \cdot 220 + 16 \cdot 150 \\ &= 2400 + 1920 + 880 + 2400 \\ &= 7600 \text{ meters} \end{aligned}$$

- (c) Bob's acceleration is  $B'(t) = 3t^2 - 12t$ .  
 $B'(5) = 3(25) - 12(5) = 15 \text{ meters/min}^2$

2 :  $\begin{cases} 1 : \text{uses } B'(t) \\ 1 : \text{answer} \end{cases}$

(d) Avg vel =  $\frac{1}{10} \int_0^{10} (t^3 - 6t^2 + 300) dt$   
 $= \frac{1}{10} \left[ \frac{t^4}{4} - 2t^3 + 300t \right]_0^{10}$   
 $= \frac{1}{10} \left[ \frac{10000}{4} - 2000 + 3000 \right] = 350 \text{ meters/min}$

3 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2016 SCORING GUIDELINES**

**Question 2**

For  $t \geq 0$ , a particle moves along the  $x$ -axis. The velocity of the particle at time  $t$  is given by

$v(t) = 1 + 2 \sin\left(\frac{t^2}{2}\right)$ . The particle is at position  $x = 2$  at time  $t = 4$ .

- (a) At time  $t = 4$ , is the particle speeding up or slowing down?  
 (b) Find all times  $t$  in the interval  $0 < t < 3$  when the particle changes direction. Justify your answer.  
 (c) Find the position of the particle at time  $t = 0$ .  
 (d) Find the total distance the particle travels from time  $t = 0$  to time  $t = 3$ .

(a)  $v(4) = 2.978716 > 0$   
 $v'(4) = -1.164000 < 0$

The particle is slowing down since the velocity and acceleration have different signs.

(b)  $v(t) = 0 \Rightarrow t = 2.707468$

$v(t)$  changes from positive to negative at  $t = 2.707$ .  
 Therefore, the particle changes direction at this time.

(c)  $x(0) = x(4) + \int_4^0 v(t) dt$   
 $= 2 + (-5.815027) = -3.815$

(d) Distance  $= \int_0^3 |v(t)| dt = 5.301$

2 : conclusion with reason

2 :  $\left\{ \begin{array}{l} 1 : t = 2.707 \\ 1 : justification \end{array} \right.$

3 :  $\left\{ \begin{array}{l} 1 : integral \\ 1 : uses initial condition \\ 1 : answer \end{array} \right.$

2 :  $\left\{ \begin{array}{l} 1 : integral \\ 1 : answer \end{array} \right.$



**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2014 SCORING GUIDELINES**

**Question 4**

Train  $A$  runs back and forth on an east-west section of railroad track. Train  $A$ 's velocity, measured in meters per minute, is given by a differentiable function  $v_A(t)$ , where time  $t$  is measured in minutes. Selected values for  $v_A(t)$  are given in the table above.

|                          |   |     |    |      |      |
|--------------------------|---|-----|----|------|------|
| $t$ (minutes)            | 0 | 2   | 5  | 8    | 12   |
| $v_A(t)$ (meters/minute) | 0 | 100 | 40 | -120 | -150 |

- (a) Find the average acceleration of train  $A$  over the interval  $2 \leq t \leq 8$ .
- (b) Do the data in the table support the conclusion that train  $A$ 's velocity is  $-100$  meters per minute at some time  $t$  with  $5 < t < 8$ ? Give a reason for your answer.
- (c) At time  $t = 2$ , train  $A$ 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train  $A$ , in meters from the Origin Station, at time  $t = 12$ . Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time  $t = 12$ .
- (d) A second train, train  $B$ , travels north from the Origin Station. At time  $t$  the velocity of train  $B$  is given by  $v_B(t) = -5t^2 + 60t + 25$ , and at time  $t = 2$  the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train  $A$  and train  $B$  is changing at time  $t = 2$ .

(a) average accel =  $\frac{v_A(8) - v_A(2)}{8 - 2} = \frac{-120 - 100}{6} = -\frac{110}{3} \text{ m/min}^2$

1 : average acceleration

(b)  $v_A$  is differentiable  $\Rightarrow v_A$  is continuous  
 $v_A(8) = -120 < -100 < 40 = v_A(5)$

2 :  $\left\{ \begin{array}{l} 1 : v_A(8) < -100 < v_A(5) \\ 1 : \text{conclusion, using IVT} \end{array} \right.$

Therefore, by the Intermediate Value Theorem, there is a time  $t$ ,  $5 < t < 8$ , such that  $v_A(t) = -100$ .

(c)  $s_A(12) = s_A(2) + \int_2^{12} v_A(t) dt = 300 + \int_2^{12} v_A(t) dt$   
 $\int_2^{12} v_A(t) dt \approx 3 \cdot \frac{100 + 40}{2} + 3 \cdot \frac{40 - 120}{2} + 4 \cdot \frac{-120 - 150}{2}$   
 $= -450$

3 :  $\left\{ \begin{array}{l} 1 : \text{position expression} \\ 1 : \text{trapezoidal sum} \\ 1 : \text{position at time } t = 12 \end{array} \right.$

$s_A(12) \approx 300 - 450 = -150$

The position of Train  $A$  at time  $t = 12$  minutes is approximately 150 meters west of Origin Station.

- (d) Let  $x$  be train  $A$ 's position,  $y$  train  $B$ 's position, and  $z$  the distance between train  $A$  and train  $B$ .

$z^2 = x^2 + y^2 \Rightarrow 2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$

$x = 300, y = 400 \Rightarrow z = 500$

$v_B(2) = -20 + 120 + 25 = 125$

$500 \frac{dz}{dt} = (300)(100) + (400)(125)$

$\frac{dz}{dt} = \frac{80000}{500} = 160 \text{ meters per minute}$

3 :  $\left\{ \begin{array}{l} 2 : \text{implicit differentiation of} \\ \quad \text{distance relationship} \\ 1 : \text{answer} \end{array} \right.$

**AP<sup>®</sup> CALCULUS AB**  
**2013 SCORING GUIDELINES**

**Question 2**

A particle moves along a straight line. For  $0 \leq t \leq 5$ , the velocity of the particle is given by

$v(t) = -2 + (t^2 + 3t)^{6/5} - t^3$ , and the position of the particle is given by  $s(t)$ . It is known that  $s(0) = 10$ .

- (a) Find all values of  $t$  in the interval  $2 \leq t \leq 4$  for which the speed of the particle is 2.
- (b) Write an expression involving an integral that gives the position  $s(t)$ . Use this expression to find the position of the particle at time  $t = 5$ .
- (c) Find all times  $t$  in the interval  $0 \leq t \leq 5$  at which the particle changes direction. Justify your answer.
- (d) Is the speed of the particle increasing or decreasing at time  $t = 4$ ? Give a reason for your answer.

- (a) Solve  $|v(t)| = 2$  on  $2 \leq t \leq 4$ .  
 $t = 3.128$  (or 3.127) and  $t = 3.473$

2 :  $\begin{cases} 1 : \text{considers } |v(t)| = 2 \\ 1 : \text{answer} \end{cases}$

(b)  $s(t) = 10 + \int_0^t v(x) dx$

$s(5) = 10 + \int_0^5 v(x) dx = -9.207$

2 :  $\begin{cases} 1 : s(t) \\ 1 : s(5) \end{cases}$

- (c)  $v(t) = 0$  when  $t = 0.536033, 3.317756$   
 $v(t)$  changes sign from negative to positive at time  $t = 0.536033$ .  
 $v(t)$  changes sign from positive to negative at time  $t = 3.317756$ .

3 :  $\begin{cases} 1 : \text{considers } v(t) = 0 \\ 2 : \text{answers with justification} \end{cases}$

Therefore, the particle changes direction at time  $t = 0.536$  and time  $t = 3.318$  (or 3.317).

- (d)  $v(4) = -11.475758 < 0$ ,  $a(4) = v'(4) = -22.295714 < 0$

2 : conclusion with reason

The speed is increasing at time  $t = 4$  because velocity and acceleration have the same sign.

**AP<sup>®</sup> CALCULUS AB  
2012 SCORING GUIDELINES**

**Question 6**

For  $0 \leq t \leq 12$ , a particle moves along the  $x$ -axis. The velocity of the particle at time  $t$  is given by

$$v(t) = \cos\left(\frac{\pi}{6}t\right). \text{ The particle is at position } x = -2 \text{ at time } t = 0.$$

- (a) For  $0 \leq t \leq 12$ , when is the particle moving to the left?
- (b) Write, but do not evaluate, an integral expression that gives the total distance traveled by the particle from time  $t = 0$  to time  $t = 6$ .
- (c) Find the acceleration of the particle at time  $t$ . Is the speed of the particle increasing, decreasing, or neither at time  $t = 4$ ? Explain your reasoning.
- (d) Find the position of the particle at time  $t = 4$ .

(a)  $v(t) = \cos\left(\frac{\pi}{6}t\right) = 0 \Rightarrow t = 3, 9$

The particle is moving to the left when  $v(t) < 0$ .  
This occurs when  $3 < t < 9$ .

(b)  $\int_0^6 |v(t)| dt$

(c)  $a(t) = -\frac{\pi}{6} \sin\left(\frac{\pi}{6}t\right)$

$$a(4) = -\frac{\pi}{6} \sin\left(\frac{2\pi}{3}\right) = -\frac{\sqrt{3}\pi}{12} < 0$$

$$v(4) = \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2} < 0$$

The speed is increasing at time  $t = 4$ , because velocity and acceleration have the same sign.

(d) 
$$\begin{aligned} x(4) &= -2 + \int_0^4 \cos\left(\frac{\pi}{6}t\right) dt \\ &= -2 + \left[\frac{6}{\pi} \sin\left(\frac{\pi}{6}t\right)\right]_0^4 \\ &= -2 + \frac{6}{\pi} \left[\sin\left(\frac{2\pi}{3}\right) - 0\right] \\ &= -2 + \frac{6}{\pi} \cdot \frac{\sqrt{3}}{2} = -2 + \frac{3\sqrt{3}}{\pi} \end{aligned}$$

2 :  $\begin{cases} 1 : \text{considers } v(t) = 0 \\ 1 : \text{interval} \end{cases}$

1 : answer

3 :  $\begin{cases} 1 : a(t) \\ 2 : \text{conclusion with reason} \end{cases}$

3 :  $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2011 SCORING GUIDELINES**

**Question 1**

For  $0 \leq t \leq 6$ , a particle is moving along the  $x$ -axis. The particle's position,  $x(t)$ , is not explicitly given. The velocity of the particle is given by  $v(t) = 2 \sin(e^{t/4}) + 1$ . The acceleration of the particle is given by  $a(t) = \frac{1}{2}e^{t/4} \cos(e^{t/4})$  and  $x(0) = 2$ .

- (a) Is the speed of the particle increasing or decreasing at time  $t = 5.5$ ? Give a reason for your answer.  
 (b) Find the average velocity of the particle for the time period  $0 \leq t \leq 6$ .  
 (c) Find the total distance traveled by the particle from time  $t = 0$  to  $t = 6$ .  
 (d) For  $0 \leq t \leq 6$ , the particle changes direction exactly once. Find the position of the particle at that time.

(a)  $v(5.5) = -0.45337$ ,  $a(5.5) = -1.35851$

The speed is increasing at time  $t = 5.5$ , because velocity and acceleration have the same sign.

2 : conclusion with reason

(b) Average velocity =  $\frac{1}{6} \int_0^6 v(t) dt = 1.949$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(c) Distance =  $\int_0^6 |v(t)| dt = 12.573$


2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(d)  $v(t) = 0$  when  $t = 5.19552$ . Let  $b = 5.19552$ .  
 $v(t)$  changes sign from positive to negative at time  $t = b$ .  
 $x(b) = 2 + \int_0^b v(t) dt = 14.134$  or  $14.135$

3 :  $\begin{cases} 1 : \text{considers } v(t) = 0 \\ 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2009 SCORING GUIDELINES**

**Question 1**



Caren rides her bicycle along a straight road from home to school, starting at home at time  $t = 0$  minutes and arriving at school at time  $t = 12$  minutes. During the time interval  $0 \leq t \leq 12$  minutes, her velocity  $v(t)$ , in miles per minute, is modeled by the piecewise-linear function whose graph is shown above.

- (a) Find the acceleration of Caren's bicycle at time  $t = 7.5$  minutes. Indicate units of measure.
- (b) Using correct units, explain the meaning of  $\int_0^{12} |v(t)| dt$  in terms of Caren's trip. Find the value of  $\int_0^{12} |v(t)| dt$ .
- (c) Shortly after leaving home, Caren realizes she left her calculus homework at home, and she returns to get it. At what time does she turn around to go back home? Give a reason for your answer.
- (d) Larry also rides his bicycle along a straight road from home to school in 12 minutes. His velocity is modeled by the function  $w$  given by  $w(t) = \frac{\pi}{15} \sin\left(\frac{\pi}{12}t\right)$ , where  $w(t)$  is in miles per minute for  $0 \leq t \leq 12$  minutes. Who lives closer to school: Caren or Larry? Show the work that leads to your answer.

(a)  $a(7.5) = v'(7.5) = \frac{v(8) - v(7)}{8 - 7} = -0.1$  miles/minute<sup>2</sup>

2 :  $\begin{cases} 1 : \text{answer} \\ 1 : \text{units} \end{cases}$

(b)  $\int_0^{12} |v(t)| dt$  is the total distance, in miles, that Caren rode during the 12 minutes from  $t = 0$  to  $t = 12$ .

2 :  $\begin{cases} 1 : \text{meaning of integral} \\ 1 : \text{value of integral} \end{cases}$

$$\int_0^{12} |v(t)| dt = \int_0^2 v(t) dt - \int_2^4 v(t) dt + \int_4^{12} v(t) dt$$

$$= 0.2 + 0.2 + 1.4 = 1.8 \text{ miles}$$

(c) Caren turns around to go back home at time  $t = 2$  minutes. This is the time at which her velocity changes from positive to negative.

2 :  $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

(d)  $\int_0^{12} w(t) dt = 1.6$ ; Larry lives 1.6 miles from school.


$$\int_0^{12} v(t) dt = 1.4; \text{ Caren lives 1.4 miles from school.}$$

Therefore, Caren lives closer to school.

3 :  $\begin{cases} 2 : \text{Larry's distance from school} \\ 1 : \text{integral} \\ 1 : \text{value} \\ 1 : \text{Caren's distance from school} \\ \text{and conclusion} \end{cases}$

**AP<sup>®</sup> CALCULUS AB  
2008 SCORING GUIDELINES**

**Question 4**



Graph of  $v$

A particle moves along the  $x$ -axis so that its velocity at time  $t$ , for  $0 \leq t \leq 6$ , is given by a differentiable function  $v$  whose graph is shown above. The velocity is 0 at  $t = 0$ ,  $t = 3$ , and  $t = 5$ , and the graph has horizontal tangents at  $t = 1$  and  $t = 4$ . The areas of the regions bounded by the  $t$ -axis and the graph of  $v$  on the intervals  $[0, 3]$ ,  $[3, 5]$ , and  $[5, 6]$  are 8, 3, and 2, respectively. At time  $t = 0$ , the particle is at  $x = -2$ .

- (a) For  $0 \leq t \leq 6$ , find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
- (b) For how many values of  $t$ , where  $0 \leq t \leq 6$ , is the particle at  $x = -8$ ? Explain your reasoning.
- (c) On the interval  $2 < t < 3$ , is the speed of the particle increasing or decreasing? Give a reason for your answer.
- (d) During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.

- (a) Since  $v(t) < 0$  for  $0 < t < 3$  and  $5 < t < 6$ , and  $v(t) > 0$  for  $3 < t < 5$ , we consider  $t = 3$  and  $t = 6$ .

$$x(3) = -2 + \int_0^3 v(t) dt = -2 - 8 = -10$$

$$x(6) = -2 + \int_0^6 v(t) dt = -2 - 8 + 3 - 2 = -9$$

Therefore, the particle is farthest left at time  $t = 3$  when its position is  $x(3) = -10$ .

- (b) The particle moves continuously and monotonically from  $x(0) = -2$  to  $x(3) = -10$ . Similarly, the particle moves continuously and monotonically from  $x(3) = -10$  to  $x(5) = -7$  and also from  $x(5) = -7$  to  $x(6) = -9$ .

By the Intermediate Value Theorem, there are three values of  $t$  for which the particle is at  $x(t) = -8$ .

- (c) The speed is decreasing on the interval  $2 < t < 3$  since on this interval  $v < 0$  and  $v$  is increasing.
- (d) The acceleration is negative on the intervals  $0 < t < 1$  and  $4 < t < 6$  since velocity is decreasing on these intervals.

$$3 : \begin{cases} 1 : \text{identifies } t = 3 \text{ as a candidate} \\ 1 : \text{considers } \int_0^6 v(t) dt \\ 1 : \text{conclusion} \end{cases}$$

$$3 : \begin{cases} 1 : \text{positions at } t = 3, t = 5, \\ \quad \text{and } t = 6 \\ 1 : \text{description of motion} \\ 1 : \text{conclusion} \end{cases}$$

1 : answer with reason

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 4**

A particle moves along the  $x$ -axis with position at time  $t$  given by  $x(t) = e^{-t} \sin t$  for  $0 \leq t \leq 2\pi$ .

- (a) Find the time  $t$  at which the particle is farthest to the left. Justify your answer.  
 (b) Find the value of the constant  $A$  for which  $x(t)$  satisfies the equation  $Ax''(t) + x'(t) + x(t) = 0$  for  $0 < t < 2\pi$ .

(a)  $x'(t) = -e^{-t} \sin t + e^{-t} \cos t = e^{-t} (\cos t - \sin t)$   
 $x'(t) = 0$  when  $\cos t = \sin t$ . Therefore,  $x'(t) = 0$  on  
 $0 \leq t \leq 2\pi$  for  $t = \frac{\pi}{4}$  and  $t = \frac{5\pi}{4}$ .

The candidates for the absolute minimum are at

$t = 0, \frac{\pi}{4}, \frac{5\pi}{4},$  and  $2\pi$ .

| $t$              | $x(t)$  |
|------------------|---|
| 0                | $e^0 \sin(0) = 0$   |
| $\frac{\pi}{4}$  | $e^{-\frac{\pi}{4}} \sin\left(\frac{\pi}{4}\right) > 0$   |
| $\frac{5\pi}{4}$ | $e^{-\frac{5\pi}{4}} \sin\left(\frac{5\pi}{4}\right) < 0$ |
| $2\pi$           | $e^{-2\pi} \sin(2\pi) = 0$                                |

The particle is farthest to the left when  $t = \frac{5\pi}{4}$ .

(b)  $x''(t) = -e^{-t} (\cos t - \sin t) + e^{-t} (-\sin t - \cos t)$   
 $= -2e^{-t} \cos t$

$$\begin{aligned} Ax''(t) + x'(t) + x(t) &= A(-2e^{-t} \cos t) + e^{-t} (\cos t - \sin t) + e^{-t} \sin t \\ &= (-2A + 1)e^{-t} \cos t \\ &= 0 \end{aligned}$$

Therefore,  $A = \frac{1}{2}$ .

$$5 : \begin{cases} 2 : x'(t) \\ 1 : \text{sets } x'(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

$$4 : \begin{cases} 2 : x''(t) \\ 1 : \text{substitutes } x''(t), x'(t), \text{ and } x(t) \\ \quad \text{into } Ax''(t) + x'(t) + x(t) \\ 1 : \text{answer} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES**

**Question 4**

|                             |   |    |    |    |    |    |    |    |    |
|-----------------------------|---|----|----|----|----|----|----|----|----|
| $t$<br>(seconds)            | 0 | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 |
| $v(t)$<br>(feet per second) | 5 | 14 | 22 | 29 | 35 | 40 | 44 | 47 | 49 |

Rocket  $A$  has positive velocity  $v(t)$  after being launched upward from an initial height of 0 feet at time  $t = 0$  seconds. The velocity of the rocket is recorded for selected values of  $t$  over the interval  $0 \leq t \leq 80$  seconds, as shown in the table above.

(a) Find the average acceleration of rocket  $A$  over the time interval  $0 \leq t \leq 80$  seconds. Indicate units of measure.

(b) Using correct units, explain the meaning of  $\int_{10}^{70} v(t) dt$  in terms of the rocket's flight. Use a midpoint

Riemann sum with 3 subintervals of equal length to approximate  $\int_{10}^{70} v(t) dt$ .

(c) Rocket  $B$  is launched upward with an acceleration of  $a(t) = \frac{3}{\sqrt{t+1}}$  feet per second per second. At time  $t = 0$  seconds, the initial height of the rocket is 0 feet, and the initial velocity is 2 feet per second. Which of the two rockets is traveling faster at time  $t = 80$  seconds? Explain your answer.

(a) Average acceleration of rocket  $A$  is

$$\frac{v(80) - v(0)}{80 - 0} = \frac{49 - 5}{80} = \frac{11}{20} \text{ ft/sec}^2$$

(b) Since the velocity is positive,  $\int_{10}^{70} v(t) dt$  represents the distance, in feet, traveled by rocket  $A$  from  $t = 10$  seconds to  $t = 70$  seconds.

A midpoint Riemann sum is

$$20[v(20) + v(40) + v(60)]$$

$$= 20[22 + 35 + 44] = 2020 \text{ ft}$$

(c) Let  $v_B(t)$  be the velocity of rocket  $B$  at time  $t$ .

$$v_B(t) = \int \frac{3}{\sqrt{t+1}} dt = 6\sqrt{t+1} + C$$

$$2 = v_B(0) = 6 + C$$

$$v_B(t) = 6\sqrt{t+1} - 4$$

$$v_B(80) = 50 > 49 = v(80)$$

Rocket  $B$  is traveling faster at time  $t = 80$  seconds.

Units of  $\text{ft/sec}^2$  in (a) and  $\text{ft}$  in (b)

1 : answer

3 :  $\left\{ \begin{array}{l} 1 : \text{explanation} \\ 1 : \text{uses } v(20), v(40), v(60) \\ 1 : \text{value} \end{array} \right.$

4 :  $\left\{ \begin{array}{l} 1 : 6\sqrt{t+1} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{finds } v_B(80), \text{ compares to } v(80), \\ \text{and draws a conclusion} \end{array} \right.$

1 : units in (a) and (b)



**AP Calculus AB**  
**Grading Rubric**  
**Integrals as**  
**Net/Accum Change**

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC**  
**2015 SCORING GUIDELINES**

**Question 1**

The rate at which rainwater flows into a drainpipe is modeled by the function  $R$ , where  $R(t) = 20 \sin\left(\frac{t^2}{35}\right)$  cubic feet per hour,  $t$  is measured in hours, and  $0 \leq t \leq 8$ . The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by  $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$  cubic feet per hour, for  $0 \leq t \leq 8$ . There are 30 cubic feet of water in the pipe at time  $t = 0$ .

- (a) How many cubic feet of rainwater flow into the pipe during the 8-hour time interval  $0 \leq t \leq 8$ ?
- (b) Is the amount of water in the pipe increasing or decreasing at time  $t = 3$  hours? Give a reason for your answer.
- (c) At what time  $t$ ,  $0 \leq t \leq 8$ , is the amount of water in the pipe at a minimum? Justify your answer.
- (d) The pipe can hold 50 cubic feet of water before overflowing. For  $t > 8$ , water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time  $w$  when the pipe will begin to overflow.

(a)  $\int_0^8 R(t) dt = 76.570$

2 :  $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b)  $R(3) - D(3) = -0.313632 < 0$   
Since  $R(3) < D(3)$ , the amount of water in the pipe is decreasing at time  $t = 3$  hours.

2 :  $\begin{cases} 1 : \text{considers } R(3) \text{ and } D(3) \\ 1 : \text{answer and reason} \end{cases}$

(c) The amount of water in the pipe at time  $t$ ,  $0 \leq t \leq 8$ , is  $30 + \int_0^t [R(x) - D(x)] dx$ .

3 :  $\begin{cases} 1 : \text{considers } R(t) - D(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

$$R(t) - D(t) = 0 \Rightarrow t = 0, 3.271658$$

| $t$      | Amount of water in the pipe |
|----------|-----------------------------|
| 0        | 30                          |
| 3.271658 | 27.964561                   |
| 8        | 48.543686                   |

The amount of water in the pipe is a minimum at time  $t = 3.272$  (or 3.271) hours.

(d)  $30 + \int_0^w [R(t) - D(t)] dt = 50$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2011 SCORING GUIDELINES**

**Question 2**

|                             |    |    |    |    |    |
|-----------------------------|----|----|----|----|----|
| $t$<br>(minutes)            | 0  | 2  | 5  | 9  | 10 |
| $H(t)$<br>(degrees Celsius) | 66 | 60 | 52 | 44 | 43 |

As a pot of tea cools, the temperature of the tea is modeled by a differentiable function  $H$  for  $0 \leq t \leq 10$ , where time  $t$  is measured in minutes and temperature  $H(t)$  is measured in degrees Celsius. Values of  $H(t)$  at selected values of time  $t$  are shown in the table above.

- (a) Use the data in the table to approximate the rate at which the temperature of the tea is changing at time  $t = 3.5$ . Show the computations that lead to your answer.
- (b) Using correct units, explain the meaning of  $\frac{1}{10} \int_0^{10} H(t) dt$  in the context of this problem. Use a trapezoidal sum with the four subintervals indicated by the table to estimate  $\frac{1}{10} \int_0^{10} H(t) dt$ .
- (c) Evaluate  $\int_0^{10} H'(t) dt$ . Using correct units, explain the meaning of the expression in the context of this problem.
- (d) At time  $t = 0$ , biscuits with temperature  $100^\circ\text{C}$  were removed from an oven. The temperature of the biscuits at time  $t$  is modeled by a differentiable function  $B$  for which it is known that  $B'(t) = -13.84e^{-0.173t}$ . Using the given models, at time  $t = 10$ , how much cooler are the biscuits than the tea?

(a) 
$$H'(3.5) \approx \frac{H(5) - H(2)}{5 - 2}$$

$$= \frac{52 - 60}{3} = -2.666 \text{ or } -2.667 \text{ degrees Celsius per minute}$$

1 : answer

- (b)  $\frac{1}{10} \int_0^{10} H(t) dt$  is the average temperature of the tea, in degrees Celsius, over the 10 minutes.

$$\frac{1}{10} \int_0^{10} H(t) dt \approx \frac{1}{10} \left( 2 \cdot \frac{66 + 60}{2} + 3 \cdot \frac{60 + 52}{2} + 4 \cdot \frac{52 + 44}{2} + 1 \cdot \frac{44 + 43}{2} \right)$$

$$= 52.95$$

3 :  $\left\{ \begin{array}{l} 1 : \text{meaning of expression} \\ 1 : \text{trapezoidal sum} \\ 1 : \text{estimate} \end{array} \right.$

(c) 
$$\int_0^{10} H'(t) dt = H(10) - H(0) = 43 - 66 = -23$$

The temperature of the tea drops 23 degrees Celsius from time  $t = 0$  to time  $t = 10$  minutes.

2 :  $\left\{ \begin{array}{l} 1 : \text{value of integral} \\ 1 : \text{meaning of expression} \end{array} \right.$

(d) 
$$B(10) = 100 + \int_0^{10} B'(t) dt = 34.18275; \quad H(10) - B(10) = 8.817$$

The biscuits are 8.817 degrees Celsius cooler than the tea.

3 :  $\left\{ \begin{array}{l} 1 : \text{integrand} \\ 1 : \text{uses } B(0) = 100 \\ 1 : \text{answer} \end{array} \right.$

**AP<sup>®</sup> CALCULUS AB**  
**2010 SCORING GUIDELINES**

**Question 1**

There is no snow on Janet's driveway when snow begins to fall at midnight. From midnight to 9 A.M., snow accumulates on the driveway at a rate modeled by  $f(t) = 7te^{\cos t}$  cubic feet per hour, where  $t$  is measured in hours since midnight. Janet starts removing snow at 6 A.M. ( $t = 6$ ). The rate  $g(t)$ , in cubic feet per hour, at which Janet removes snow from the driveway at time  $t$  hours after midnight is modeled by

$$g(t) = \begin{cases} 0 & \text{for } 0 \leq t < 6 \\ 125 & \text{for } 6 \leq t < 7 \\ 108 & \text{for } 7 \leq t \leq 9. \end{cases}$$

- (a) How many cubic feet of snow have accumulated on the driveway by 6 A.M.?  
 (b) Find the rate of change of the volume of snow on the driveway at 8 A.M.  
 (c) Let  $h(t)$  represent the total amount of snow, in cubic feet, that Janet has removed from the driveway at time  $t$  hours after midnight. Express  $h$  as a piecewise-defined function with domain  $0 \leq t \leq 9$ .  
 (d) How many cubic feet of snow are on the driveway at 9 A.M.?

(a)  $\int_0^6 f(t) dt = 142.274$  or  $142.275$  cubic feet

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) Rate of change is  $f(8) - g(8) = -59.582$  or  $-59.583$  cubic feet per hour.

1 : answer

(c)  $h(0) = 0$

For  $0 < t \leq 6$ ,  $h(t) = h(0) + \int_0^t g(s) ds = 0 + \int_0^t 0 ds = 0$ .

For  $6 < t \leq 7$ ,  $h(t) = h(6) + \int_6^t g(s) ds = 0 + \int_6^t 125 ds = 125(t - 6)$ .

For  $7 < t \leq 9$ ,  $h(t) = h(7) + \int_7^t g(s) ds = 125 + \int_7^t 108 ds = 125 + 108(t - 7)$ .

Thus,  $h(t) = \begin{cases} 0 & \text{for } 0 \leq t \leq 6 \\ 125(t - 6) & \text{for } 6 < t \leq 7 \\ 125 + 108(t - 7) & \text{for } 7 < t \leq 9 \end{cases}$

3 :  $\begin{cases} 1 : h(t) \text{ for } 0 \leq t \leq 6 \\ 1 : h(t) \text{ for } 6 < t \leq 7 \\ 1 : h(t) \text{ for } 7 < t \leq 9 \end{cases}$


(d) Amount of snow is  $\int_0^9 f(t) dt - h(9) = 26.334$  or  $26.335$  cubic feet.

3 :  $\begin{cases} 1 : \text{integral} \\ 1 : h(9) \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2010 SCORING GUIDELINES**

**Question 3**

There are 700 people in line for a popular amusement-park ride when the ride begins operation in the morning. Once it begins operation, the ride accepts passengers until the park closes 8 hours later. While there is a line, people move onto the ride at a rate of 800 people per hour. The graph above shows the rate,  $r(t)$ , at which people arrive at the ride throughout the day. Time  $t$  is measured in hours from the time the ride begins operation.



- (a) How many people arrive at the ride between  $t = 0$  and  $t = 3$ ? Show the computations that lead to your answer.
- (b) Is the number of people waiting in line to get on the ride increasing or decreasing between  $t = 2$  and  $t = 3$ ? Justify your answer.
- (c) At what time  $t$  is the line for the ride the longest? How many people are in line at that time? Justify your answers.
- (d) Write, but do not solve, an equation involving an integral expression of  $r$  whose solution gives the earliest time  $t$  at which there is no longer a line for the ride.

(a)  $\int_0^3 r(t) dt = 2 \cdot \frac{1000 + 1200}{2} + \frac{1200 + 800}{2} = 3200$  people

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) The number of people waiting in line is increasing because people move onto the ride at a rate of 800 people per hour and for  $2 < t < 3$ ,  $r(t) > 800$ .

1 : answer with reason

(c)  $r(t) = 800$  only at  $t = 3$   
For  $0 \leq t < 3$ ,  $r(t) > 800$ . For  $3 < t \leq 8$ ,  $r(t) < 800$ .  
Therefore, the line is longest at time  $t = 3$ .  
There are  $700 + 3200 - 800 \cdot 3 = 1500$  people waiting in line at time  $t = 3$ .

3 :  $\begin{cases} 1 : \text{identifies } t = 3 \\ 1 : \text{number of people in line} \\ 1 : \text{justification} \end{cases}$

(d)  $0 = 700 + \int_0^t r(s) ds - 800t$

3 :  $\begin{cases} 1 : 800t \\ 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2009 SCORING GUIDELINES**

**Question 2**

The rate at which people enter an auditorium for a rock concert is modeled by the function  $R$  given by  $R(t) = 1380t^2 - 675t^3$  for  $0 \leq t \leq 2$  hours;  $R(t)$  is measured in people per hour. No one is in the auditorium at time  $t = 0$ , when the doors open. The doors close and the concert begins at time  $t = 2$ .

- (a) How many people are in the auditorium when the concert begins?
- (b) Find the time when the rate at which people enter the auditorium is a maximum. Justify your answer.
- (c) The total wait time for all the people in the auditorium is found by adding the time each person waits, starting at the time the person enters the auditorium and ending when the concert begins. The function  $w$  models the total wait time for all the people who enter the auditorium before time  $t$ . The derivative of  $w$  is given by  $w'(t) = (2 - t)R(t)$ . Find  $w(2) - w(1)$ , the total wait time for those who enter the auditorium after time  $t = 1$ .
- (d) On average, how long does a person wait in the auditorium for the concert to begin? Consider all people who enter the auditorium after the doors open, and use the model for total wait time from part (c).

(a)  $\int_0^2 R(t) dt = 980$  people

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b)  $R'(t) = 0$  when  $t = 0$  and  $t = 1.36296$   
The maximum rate may occur at 0,  $a = 1.36296$ , or 2.

$$R(0) = 0$$

$$R(a) = 854.527$$

$$R(2) = 120$$

The maximum rate occurs when  $t = 1.362$  or  $1.363$ .

3 :  $\begin{cases} 1 : \text{considers } R'(t) = 0 \\ 1 : \text{interior critical point} \\ 1 : \text{answer and justification} \end{cases}$

(c)  $w(2) - w(1) = \int_1^2 w'(t) dt = \int_1^2 (2 - t)R(t) dt = 387.5$

The total wait time for those who enter the auditorium after time  $t = 1$  is 387.5 hours.

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(d)  $\frac{1}{980}w(2) = \frac{1}{980} \int_0^2 (2 - t)R(t) dt = 0.77551$

On average, a person waits 0.775 or 0.776 hour.

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2009 SCORING GUIDELINES**

**Question 3**

Mighty Cable Company manufactures cable that sells for \$120 per meter. For a cable of fixed length, the cost of producing a portion of the cable varies with its distance from the beginning of the cable. Mighty reports that the cost to produce a portion of a cable that is  $x$  meters from the beginning of the cable is  $6\sqrt{x}$  dollars per meter. (Note: Profit is defined to be the difference between the amount of money received by the company for selling the cable and the company's cost of producing the cable.)

- (a) Find Mighty's profit on the sale of a 25-meter cable.
- (b) Using correct units, explain the meaning of  $\int_{25}^{30} 6\sqrt{x} \, dx$  in the context of this problem.
- (c) Write an expression, involving an integral, that represents Mighty's profit on the sale of a cable that is  $k$  meters long.
- (d) Find the maximum profit that Mighty could earn on the sale of one cable. Justify your answer.

(a) Profit =  $120 \cdot 25 - \int_0^{25} 6\sqrt{x} \, dx = 2500$  dollars

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b)  $\int_{25}^{30} 6\sqrt{x} \, dx$  is the difference in cost, in dollars, of producing a cable of length 30 meters and a cable of length 25 meters.

1 : answer with units

(c) Profit =  $120k - \int_0^k 6\sqrt{x} \, dx$  dollars

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{expression} \end{cases}$

(d) Let  $P(k)$  be the profit for a cable of length  $k$ .

$$P'(k) = 120 - 6\sqrt{k} = 0 \text{ when } k = 400.$$

This is the only critical point for  $P$ , and  $P'$  changes from positive to negative at  $k = 400$ .

Therefore, the maximum profit is  $P(400) = 16,000$  dollars.

4 :  $\begin{cases} 1 : P'(k) = 0 \\ 1 : k = 400 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2008 SCORING GUIDELINES**

**Question 2**

|                 |     |     |     |     |     |    |   |
|-----------------|-----|-----|-----|-----|-----|----|---|
| $t$ (hours)     | 0   | 1   | 3   | 4   | 7   | 8  | 9 |
| $L(t)$ (people) | 120 | 156 | 176 | 126 | 150 | 80 | 0 |

Concert tickets went on sale at noon ( $t = 0$ ) and were sold out within 9 hours. The number of people waiting in line to purchase tickets at time  $t$  is modeled by a twice-differentiable function  $L$  for  $0 \leq t \leq 9$ . Values of  $L(t)$  at various times  $t$  are shown in the table above.

- (a) Use the data in the table to estimate the rate at which the number of people waiting in line was changing at 5:30 P.M. ( $t = 5.5$ ). Show the computations that lead to your answer. Indicate units of measure.
- (b) Use a trapezoidal sum with three subintervals to estimate the average number of people waiting in line during the first 4 hours that tickets were on sale.
- (c) For  $0 \leq t \leq 9$ , what is the fewest number of times at which  $L'(t)$  must equal 0? Give a reason for your answer.
- (d) The rate at which tickets were sold for  $0 \leq t \leq 9$  is modeled by  $r(t) = 550te^{-t/2}$  tickets per hour. Based on the model, how many tickets were sold by 3 P.M. ( $t = 3$ ), to the nearest whole number?

(a)  $L'(5.5) \approx \frac{L(7) - L(4)}{7 - 4} = \frac{150 - 126}{3} = 8$  people per hour

2 : { 1 : estimate  
1 : units

(b) The average number of people waiting in line during the first 4 hours is approximately

$$\frac{1}{4} \left( \frac{L(0) + L(1)}{2} (1 - 0) + \frac{L(1) + L(3)}{2} (3 - 1) + \frac{L(3) + L(4)}{2} (4 - 3) \right)$$

$= 155.25$  people

2 : { 1 : trapezoidal sum  
1 : answer

(c)  $L$  is differentiable on  $[0, 9]$  so the Mean Value Theorem implies  $L'(t) > 0$  for some  $t$  in  $(1, 3)$  and some  $t$  in  $(4, 7)$ . Similarly,  $L'(t) < 0$  for some  $t$  in  $(3, 4)$  and some  $t$  in  $(7, 8)$ . Then, since  $L'$  is continuous on  $[0, 9]$ , the Intermediate Value Theorem implies that  $L'(t) = 0$  for at least three values of  $t$  in  $[0, 9]$ .

3 : { 1 : considers change in sign of  $L'$   
1 : analysis  
1 : conclusion

OR

The continuity of  $L$  on  $[1, 4]$  implies that  $L$  attains a maximum value there. Since  $L(3) > L(1)$  and  $L(3) > L(4)$ , this maximum occurs on  $(1, 4)$ . Similarly,  $L$  attains a minimum on  $(3, 7)$  and a maximum on  $(4, 8)$ .  $L$  is differentiable, so  $L'(t) = 0$  at each relative extreme point on  $(0, 9)$ . Therefore  $L'(t) = 0$  for at least three values of  $t$  in  $[0, 9]$ .

OR

3 : { 1 : considers relative extrema of  $L$  on  $(0, 9)$   
1 : analysis  
1 : conclusion

[Note: There is a function  $L$  that satisfies the given conditions with  $L'(t) = 0$  for exactly three values of  $t$ .]

(d)  $\int_0^3 r(t) dt = 972.784$

There were approximately 973 tickets sold by 3 P.M.

2 : { 1 : integrand  
1 : limits and answer



**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 2**


The amount of water in a storage tank, in gallons, is modeled by a continuous function on the time interval  $0 \leq t \leq 7$ , where  $t$  is measured in hours. In this model, rates are given as follows:

(i) The rate at which water enters the tank is

$$f(t) = 100t^2 \sin(\sqrt{t}) \text{ gallons per hour for } 0 \leq t \leq 7.$$

(ii) The rate at which water leaves the tank is

$$g(t) = \begin{cases} 250 & \text{for } 0 \leq t < 3 \\ 2000 & \text{for } 3 < t \leq 7 \end{cases} \text{ gallons per hour.}$$



The graphs of  $f$  and  $g$ , which intersect at  $t = 1.617$  and  $t = 5.076$ , are shown in the figure above. At time  $t = 0$ , the amount of water in the tank is 5000 gallons.

- (a) How many gallons of water enter the tank during the time interval  $0 \leq t \leq 7$ ? Round your answer to the nearest gallon.
- (b) For  $0 \leq t \leq 7$ , find the time intervals during which the amount of water in the tank is decreasing. Give a reason for each answer.
- (c) For  $0 \leq t \leq 7$ , at what time  $t$  is the amount of water in the tank greatest? To the nearest gallon, compute the amount of water at this time. Justify your answer.

(a)  $\int_0^7 f(t) dt \approx 8264$  gallons

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) The amount of water in the tank is decreasing on the intervals  $0 \leq t \leq 1.617$  and  $3 \leq t \leq 5.076$  because  $f(t) < g(t)$  for  $0 \leq t < 1.617$  and  $3 < t < 5.076$ .

2 :  $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(c) Since  $f(t) - g(t)$  changes sign from positive to negative only at  $t = 3$ , the candidates for the absolute maximum are at  $t = 0, 3$ , and  $7$ .

5 :  $\begin{cases} 1 : \text{identifies } t = 3 \text{ as a candidate} \\ 1 : \text{integrand} \\ 1 : \text{amount of water at } t = 3 \\ 1 : \text{amount of water at } t = 7 \\ 1 : \text{conclusion} \end{cases}$


| $t$ (hours) | gallons of water                                   |
|-------------|--|
| 0           | 5000   |
| 3           | $5000 + \int_0^3 f(t) dt - 250(3) = 5126.591$      |
| 7           | $5126.591 + \int_3^7 f(t) dt - 2000(4) = 4513.807$ |

The amount of water in the tank is greatest at 3 hours. At that time, the amount of water in the tank, rounded to the nearest gallon, is 5127 gallons.

**AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES**

**Question 2**

At an intersection in Thomasville, Oregon, cars turn left at the rate  $L(t) = 60\sqrt{t} \sin^2\left(\frac{t}{3}\right)$  cars per hour over the time interval  $0 \leq t \leq 18$  hours. The graph of  $y = L(t)$  is shown above.



- (a) To the nearest whole number, find the total number of cars turning left at the intersection over the time interval  $0 \leq t \leq 18$  hours.
- (b) Traffic engineers will consider turn restrictions when  $L(t) \geq 150$  cars per hour. Find all values of  $t$  for which  $L(t) \geq 150$  and compute the average value of  $L$  over this time interval. Indicate units of measure.
- (c) Traffic engineers will install a signal if there is any two-hour time interval during which the product of the total number of cars turning left and the total number of oncoming cars traveling straight through the intersection is greater than 200,000. In every two-hour time interval, 500 oncoming cars travel straight through the intersection. Does this intersection require a traffic signal? Explain the reasoning that leads to your conclusion.

(a)  $\int_0^{18} L(t) dt \approx 1658$  cars

2 :  $\left\{ \begin{array}{l} 1 : \text{setup} \\ 1 : \text{answer} \end{array} \right.$

(b)  $L(t) = 150$  when  $t = 12.42831, 16.12166$

Let  $R = 12.42831$  and  $S = 16.12166$

$L(t) \geq 150$  for  $t$  in the interval  $[R, S]$

$\frac{1}{S-R} \int_R^S L(t) dt = 199.426$  cars per hour

3 :  $\left\{ \begin{array}{l} 1 : t\text{-interval when } L(t) \geq 150 \\ 1 : \text{average value integral} \\ 1 : \text{answer with units} \end{array} \right.$

- (c) For the product to exceed 200,000, the number of cars turning left in a two-hour interval must be greater than 400.

$\int_{13}^{15} L(t) dt = 431.931 > 400$

4 :  $\left\{ \begin{array}{l} 1 : \text{considers 400 cars} \\ 1 : \text{valid interval } [h, h+2] \\ 1 : \text{value of } \int_h^{h+2} L(t) dt \\ 1 : \text{answer and explanation} \end{array} \right.$

OR

The number of cars turning left will be greater than 400 on a two-hour interval if  $L(t) \geq 200$  on that interval.

$L(t) \geq 200$  on any two-hour subinterval of  $[13.25304, 15.32386]$ .

OR

4 :  $\left\{ \begin{array}{l} 1 : \text{considers 200 cars per hour} \\ 1 : \text{solves } L(t) \geq 200 \\ 1 : \text{discusses 2 hour interval} \\ 1 : \text{answer and explanation} \end{array} \right.$

Yes, a traffic signal is required.

**AP Calculus AB**  
**Grading Rubric**  
**Riemann Sums, IVT,**  
**EVT, and MVT**

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2016 SCORING GUIDELINES**

**Question 1**

|                           |      |      |     |     |     |
|---------------------------|------|------|-----|-----|-----|
| $t$<br>(hours)            | 0    | 1    | 3   | 6   | 8   |
| $R(t)$<br>(liters / hour) | 1340 | 1190 | 950 | 740 | 700 |

Water is pumped into a tank at a rate modeled by  $W(t) = 2000e^{-t^2/20}$  liters per hour for  $0 \leq t \leq 8$ , where  $t$  is measured in hours. Water is removed from the tank at a rate modeled by  $R(t)$  liters per hour, where  $R$  is differentiable and decreasing on  $0 \leq t \leq 8$ . Selected values of  $R(t)$  are shown in the table above. At time  $t = 0$ , there are 50,000 liters of water in the tank.

- (a) Estimate  $R'(2)$ . Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For  $0 \leq t \leq 8$ , is there a time  $t$  when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

(a)  $R'(2) \approx \frac{R(3) - R(1)}{3 - 1} = \frac{950 - 1190}{3 - 1} = -120 \text{ liters/hr}^2$

2 :  $\begin{cases} 1 : \text{estimate} \\ 1 : \text{units} \end{cases}$

(b) The total amount of water removed is given by  $\int_0^8 R(t) dt$ .

$$\begin{aligned} \int_0^8 R(t) dt &\approx 1 \cdot R(0) + 2 \cdot R(1) + 3 \cdot R(3) + 2 \cdot R(6) \\ &= 1(1340) + 2(1190) + 3(950) + 2(740) \\ &= 8050 \text{ liters} \end{aligned}$$

This is an overestimate since  $R$  is a decreasing function.

3 :  $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{estimate} \\ 1 : \text{overestimate with reason} \end{cases}$

(c)  $\text{Total} \approx 50000 + \int_0^8 W(t) dt - 8050$   
 $= 50000 + 7836.195325 - 8050 \approx 49786 \text{ liters}$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{estimate} \end{cases}$

(d)  $W(0) - R(0) > 0$ ,  $W(8) - R(8) < 0$ , and  $W(t) - R(t)$  is continuous.

2 :  $\begin{cases} 1 : \text{considers } W(t) - R(t) \\ 1 : \text{answer with explanation} \end{cases}$

Therefore, the Intermediate Value Theorem guarantees at least one time  $t$ ,  $0 < t < 8$ , for which  $W(t) - R(t) = 0$ , or  $W(t) = R(t)$ .

For this value of  $t$ , the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank.

**AP<sup>®</sup> CALCULUS AB**  
**2013 SCORING GUIDELINES**

**Question 3**

|                    |   |     |     |      |      |      |      |
|--------------------|---|-----|-----|------|------|------|------|
| $t$<br>(minutes)   | 0 | 1   | 2   | 3    | 4    | 5    | 6    |
| $C(t)$<br>(ounces) | 0 | 5.3 | 8.8 | 11.2 | 12.8 | 13.8 | 14.5 |

Hot water is dripping through a coffeemaker, filling a large cup with coffee. The amount of coffee in the cup at time  $t$ ,  $0 \leq t \leq 6$ , is given by a differentiable function  $C$ , where  $t$  is measured in minutes. Selected values of  $C(t)$ , measured in ounces, are given in the table above.

- (a) Use the data in the table to approximate  $C'(3.5)$ . Show the computations that lead to your answer, and indicate units of measure.
- (b) Is there a time  $t$ ,  $2 \leq t \leq 4$ , at which  $C'(t) = 2$ ? Justify your answer.
- (c) Use a midpoint sum with three subintervals of equal length indicated by the data in the table to approximate the value of  $\frac{1}{6} \int_0^6 C(t) dt$ . Using correct units, explain the meaning of  $\frac{1}{6} \int_0^6 C(t) dt$  in the context of the problem.
- (d) The amount of coffee in the cup, in ounces, is modeled by  $B(t) = 16 - 16e^{-0.4t}$ . Using this model, find the rate at which the amount of coffee in the cup is changing when  $t = 5$ .

(a)  $C'(3.5) \approx \frac{C(4) - C(3)}{4 - 3} = \frac{12.8 - 11.2}{1} = 1.6$  ounces/min

2 :  $\begin{cases} 1 : \text{approximation} \\ 1 : \text{units} \end{cases}$

(b)  $C$  is differentiable  $\Rightarrow C$  is continuous (on the closed interval)

$$\frac{C(4) - C(2)}{4 - 2} = \frac{12.8 - 8.8}{2} = 2$$

Therefore, by the Mean Value Theorem, there is at least one time  $t$ ,  $2 < t < 4$ , for which  $C'(t) = 2$ .

2 :  $\begin{cases} 1 : \frac{C(4) - C(2)}{4 - 2} \\ 1 : \text{conclusion, using MVT} \end{cases}$

(c)  $\frac{1}{6} \int_0^6 C(t) dt \approx \frac{1}{6} [2 \cdot C(1) + 2 \cdot C(3) + 2 \cdot C(5)]$   
 $= \frac{1}{6} (2 \cdot 5.3 + 2 \cdot 11.2 + 2 \cdot 13.8)$   
 $= \frac{1}{6} (60.6) = 10.1$  ounces

3 :  $\begin{cases} 1 : \text{midpoint sum} \\ 1 : \text{approximation} \\ 1 : \text{interpretation} \end{cases}$

$\frac{1}{6} \int_0^6 C(t) dt$  is the average amount of coffee in the cup, in ounces, over the time interval  $0 \leq t \leq 6$  minutes.

(d)  $B'(t) = -16(-0.4)e^{-0.4t} = 6.4e^{-0.4t}$   
 $B'(5) = 6.4e^{-0.4(5)} = \frac{6.4}{e^2}$  ounces/min

2 :  $\begin{cases} 1 : B'(t) \\ 1 : B'(5) \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2012 SCORING GUIDELINES**

**Question 1**

|                             |      |      |      |      |      |
|-----------------------------|------|------|------|------|------|
| $t$ (minutes)               | 0    | 4    | 9    | 15   | 20   |
| $W(t)$ (degrees Fahrenheit) | 55.0 | 57.1 | 61.8 | 67.9 | 71.0 |

The temperature of water in a tub at time  $t$  is modeled by a strictly increasing, twice-differentiable function  $W$ , where  $W(t)$  is measured in degrees Fahrenheit and  $t$  is measured in minutes. At time  $t = 0$ , the temperature of the water is  $55^\circ\text{F}$ . The water is heated for 30 minutes, beginning at time  $t = 0$ . Values of  $W(t)$  at selected times  $t$  for the first 20 minutes are given in the table above.

- (a) Use the data in the table to estimate  $W'(12)$ . Show the computations that lead to your answer. Using correct units, interpret the meaning of your answer in the context of this problem.
- (b) Use the data in the table to evaluate  $\int_0^{20} W'(t) dt$ . Using correct units, interpret the meaning of  $\int_0^{20} W'(t) dt$  in the context of this problem.
- (c) For  $0 \leq t \leq 20$ , the average temperature of the water in the tub is  $\frac{1}{20} \int_0^{20} W(t) dt$ . Use a left Riemann sum with the four subintervals indicated by the data in the table to approximate  $\frac{1}{20} \int_0^{20} W(t) dt$ . Does this approximation overestimate or underestimate the average temperature of the water over these 20 minutes? Explain your reasoning.
- (d) For  $20 \leq t \leq 25$ , the function  $W$  that models the water temperature has first derivative given by  $W'(t) = 0.4\sqrt{t} \cos(0.06t)$ . Based on the model, what is the temperature of the water at time  $t = 25$ ?

(a) 
$$W'(12) \approx \frac{W(15) - W(9)}{15 - 9} = \frac{67.9 - 61.8}{6}$$

$$= 1.017 \text{ (or } 1.016)$$

The water temperature is increasing at a rate of approximately  $1.017^\circ\text{F}$  per minute at time  $t = 12$  minutes.

(b) 
$$\int_0^{20} W'(t) dt = W(20) - W(0) = 71.0 - 55.0 = 16$$

The water has warmed by  $16^\circ\text{F}$  over the interval from  $t = 0$  to  $t = 20$  minutes.

(c) 
$$\frac{1}{20} \int_0^{20} W(t) dt \approx \frac{1}{20} (4 \cdot W(0) + 5 \cdot W(4) + 6 \cdot W(9) + 5 \cdot W(15))$$

$$= \frac{1}{20} (4 \cdot 55.0 + 5 \cdot 57.1 + 6 \cdot 61.8 + 5 \cdot 67.9)$$

$$= \frac{1}{20} \cdot 1215.8 = 60.79$$

This approximation is an underestimate, because a left Riemann sum is used and the function  $W$  is strictly increasing.

(d) 
$$W(25) = 71.0 + \int_{20}^{25} W'(t) dt$$

$$= 71.0 + 2.043155 = 73.043$$

2 :  $\begin{cases} 1 : \text{estimate} \\ 1 : \text{interpretation with units} \end{cases}$

2 :  $\begin{cases} 1 : \text{value} \\ 1 : \text{interpretation with units} \end{cases}$

3 :  $\begin{cases} 1 : \text{left Riemann sum} \\ 1 : \text{approximation} \\ 1 : \text{underestimate with reason} \end{cases}$

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2010 SCORING GUIDELINES**

**Question 2**

|                                 |   |   |    |    |    |
|---------------------------------|---|---|----|----|----|
| $t$<br>(hours)                  | 0 | 2 | 5  | 7  | 8  |
| $E(t)$<br>(hundreds of entries) | 0 | 4 | 13 | 21 | 23 |

A zoo sponsored a one-day contest to name a new baby elephant. Zoo visitors deposited entries in a special box between noon ( $t = 0$ ) and 8 P.M. ( $t = 8$ ). The number of entries in the box  $t$  hours after noon is modeled by a differentiable function  $E$  for  $0 \leq t \leq 8$ . Values of  $E(t)$ , in hundreds of entries, at various times  $t$  are shown in the table above.

(a) Use the data in the table to approximate the rate, in hundreds of entries per hour, at which entries were being deposited at time  $t = 6$ . Show the computations that lead to your answer.

(b) Use a trapezoidal sum with the four subintervals given by the table to approximate the value of  $\frac{1}{8} \int_0^8 E(t) dt$ .

Using correct units, explain the meaning of  $\frac{1}{8} \int_0^8 E(t) dt$  in terms of the number of entries.

(c) At 8 P.M., volunteers began to process the entries. They processed the entries at a rate modeled by the function  $P$ , where  $P(t) = t^3 - 30t^2 + 298t - 976$  hundreds of entries per hour for  $8 \leq t \leq 12$ . According to the model, how many entries had not yet been processed by midnight ( $t = 12$ )?

(d) According to the model from part (c), at what time were the entries being processed most quickly? Justify your answer.

(a)  $E'(6) \approx \frac{E(7) - E(5)}{7 - 5} = 4$  hundred entries per hour

(b)  $\frac{1}{8} \int_0^8 E(t) dt \approx$   
 $\frac{1}{8} \left( 2 \cdot \frac{E(0) + E(2)}{2} + 3 \cdot \frac{E(2) + E(5)}{2} + 2 \cdot \frac{E(5) + E(7)}{2} + 1 \cdot \frac{E(7) + E(8)}{2} \right)$   
 $= 10.687$  or  $10.688$

$\frac{1}{8} \int_0^8 E(t) dt$  is the average number of hundreds of entries in the box between noon and 8 P.M.

(c)  $23 - \int_8^{12} P(t) dt = 23 - 16 = 7$  hundred entries

(d)  $P'(t) = 0$  when  $t = 9.183503$  and  $t = 10.816497$ .

|           |          |
|-----------|----------|
| $t$       | $P(t)$   |
| 8         | 0        |
| 9.183503  | 5.088662 |
| 10.816497 | 2.911338 |
| 12        | 8        |

Entries are being processed most quickly at time  $t = 12$ .

1 : answer

3 :  $\left\{ \begin{array}{l} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \\ 1 : \text{meaning} \end{array} \right.$

2 :  $\left\{ \begin{array}{l} 1 : \text{integral} \\ 1 : \text{answer} \end{array} \right.$

3 :  $\left\{ \begin{array}{l} 1 : \text{considers } P'(t) = 0 \\ 1 : \text{identifies candidates} \\ 1 : \text{answer with justification} \end{array} \right.$

**AP<sup>®</sup> CALCULUS AB**  
**2009 SCORING GUIDELINES**

**Question 5**

|        |   |   |    |   |    |
|--------|---|---|----|---|----|
| $x$    | 2 | 3 | 5  | 8 | 13 |
| $f(x)$ | 1 | 4 | -2 | 3 | 6  |

Let  $f$  be a function that is twice differentiable for all real numbers. The table above gives values of  $f$  for selected points in the closed interval  $2 \leq x \leq 13$ .

- (a) Estimate  $f'(4)$ . Show the work that leads to your answer.
- (b) Evaluate  $\int_2^{13} (3 - 5f''(x)) dx$ . Show the work that leads to your answer.
- (c) Use a left Riemann sum with subintervals indicated by the data in the table to approximate  $\int_2^{13} f(x) dx$ . Show the work that leads to your answer.
- (d) Suppose  $f'(5) = 3$  and  $f''(x) < 0$  for all  $x$  in the closed interval  $5 \leq x \leq 8$ . Use the line tangent to the graph of  $f$  at  $x = 5$  to show that  $f(7) \leq 4$ . Use the secant line for the graph of  $f$  on  $5 \leq x \leq 8$  to show that  $f(7) \geq \frac{4}{3}$ .

(a)  $f'(4) \approx \frac{f(5) - f(3)}{5 - 3} = -3$

(b) 
$$\int_2^{13} (3 - 5f''(x)) dx = \int_2^{13} 3 dx - 5 \int_2^{13} f''(x) dx$$

$$= 3(13 - 2) - 5(f'(13) - f'(2)) = 8$$

(c) 
$$\int_2^{13} f(x) dx \approx f(2)(3 - 2) + f(3)(5 - 3)$$

$$+ f(5)(8 - 5) + f(8)(13 - 8) = 18$$

- (d) An equation for the tangent line is  $y = -2 + 3(x - 5)$ .  
 Since  $f''(x) < 0$  for all  $x$  in the interval  $5 \leq x \leq 8$ , the line tangent to the graph of  $y = f(x)$  at  $x = 5$  lies above the graph for all  $x$  in the interval  $5 < x \leq 8$ .  
 Therefore,  $f(7) \leq -2 + 3 \cdot 2 = 4$ .

An equation for the secant line is  $y = -2 + \frac{5}{3}(x - 5)$ .

Since  $f''(x) < 0$  for all  $x$  in the interval  $5 \leq x \leq 8$ , the secant line connecting  $(5, f(5))$  and  $(8, f(8))$  lies below the graph of  $y = f(x)$  for all  $x$  in the interval  $5 < x < 8$ .

Therefore,  $f(7) \geq -2 + \frac{5}{3} \cdot 2 = \frac{4}{3}$ .

1 : answer

2 :  $\left\{ \begin{array}{l} 1 : \text{uses Fundamental Theorem} \\ \quad \text{of Calculus} \\ 1 : \text{answer} \end{array} \right.$

2 :  $\left\{ \begin{array}{l} 1 : \text{left Riemann sum} \\ 1 : \text{answer} \end{array} \right.$

4 :  $\left\{ \begin{array}{l} 1 : \text{tangent line} \\ 1 : \text{shows } f(7) \leq 4 \\ 1 : \text{secant line} \\ 1 : \text{shows } f(7) \geq \frac{4}{3} \end{array} \right.$



**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 3**

| $x$ | $f(x)$ | $f'(x)$ | $g(x)$ | $g'(x)$ |
|-----|--------|---------|--------|---------|
| 1   | 6      | 4       | 2      | 5       |
| 2   | 9      | 2       | 3      | 1       |
| 3   | 10     | -4      | 4      | 2       |
| 4   | -1     | 3       | 6      | 7       |

The functions  $f$  and  $g$  are differentiable for all real numbers, and  $g$  is strictly increasing. The table above gives values of the functions and their first derivatives at selected values of  $x$ . The function  $h$  is given by  $h(x) = f(g(x)) - 6$ .

- (a) Explain why there must be a value  $r$  for  $1 < r < 3$  such that  $h(r) = -5$ .
- (b) Explain why there must be a value  $c$  for  $1 < c < 3$  such that  $h'(c) = -5$ .
- (c) Let  $w$  be the function given by  $w(x) = \int_1^{g(x)} f(t) dt$ . Find the value of  $w'(3)$ .
- (d) If  $g^{-1}$  is the inverse function of  $g$ , write an equation for the line tangent to the graph of  $y = g^{-1}(x)$  at  $x = 2$ .

(a)  $h(1) = f(g(1)) - 6 = f(2) - 6 = 9 - 6 = 3$   
 $h(3) = f(g(3)) - 6 = f(4) - 6 = -1 - 6 = -7$   
 Since  $h(3) < -5 < h(1)$  and  $h$  is continuous, by the Intermediate Value Theorem, there exists a value  $r$ ,  $1 < r < 3$ , such that  $h(r) = -5$ .

(b)  $\frac{h(3) - h(1)}{3 - 1} = \frac{-7 - 3}{3 - 1} = -5$   
 Since  $h$  is continuous and differentiable, by the Mean Value Theorem, there exists a value  $c$ ,  $1 < c < 3$ , such that  $h'(c) = -5$ .

(c)  $w'(3) = f(g(3)) \cdot g'(3) = f(4) \cdot 2 = -2$

(d)  $g(1) = 2$ , so  $g^{-1}(2) = 1$ .  
 $(g^{-1})'(2) = \frac{1}{g'(g^{-1}(2))} = \frac{1}{g'(1)} = \frac{1}{5}$   
 An equation of the tangent line is  $y - 1 = \frac{1}{5}(x - 2)$ .

2 :  $\begin{cases} 1 : h(1) \text{ and } h(3) \\ 1 : \text{conclusion, using IVT} \end{cases}$

2 :  $\begin{cases} 1 : \frac{h(3) - h(1)}{3 - 1} \\ 1 : \text{conclusion, using MVT} \end{cases}$

2 :  $\begin{cases} 1 : \text{apply chain rule} \\ 1 : \text{answer} \end{cases}$

3 :  $\begin{cases} 1 : g^{-1}(2) \\ 1 : (g^{-1})'(2) \\ 1 : \text{tangent line equation} \end{cases}$


**AP Calculus AB**  
**Grading Rubric**  
**Differential EQS, Slope**  
**Fields, and Related**  
**Rates**

**AP<sup>®</sup> CALCULUS AB**  
**2016 SCORING GUIDELINES**

**Question 4**

Consider the differential equation  $\frac{dy}{dx} = \frac{y^2}{x-1}$ .

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Let  $y = f(x)$  be the particular solution to the given differential equation with the initial condition  $f(2) = 3$ . Write an equation for the line tangent to the graph of  $y = f(x)$  at  $x = 2$ .  
Use your equation to approximate  $f(2.1)$ .
- (c) Find the particular solution  $y = f(x)$  to the given differential equation with the initial condition  $f(2) = 3$ .



(b)  $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = \frac{3^2}{2-1} = 9$

An equation for the tangent line is  $y = 9(x - 2) + 3$ .

$$f(2.1) \approx 9(2.1 - 2) + 3 = 3.9$$

(c)  $\frac{1}{y^2} dy = \frac{1}{x-1} dx$

$$\int \frac{1}{y^2} dy = \int \frac{1}{x-1} dx$$

$$-\frac{1}{y} = \ln|x-1| + C$$

$$-\frac{1}{3} = \ln|2-1| + C \Rightarrow C = -\frac{1}{3}$$

$$-\frac{1}{y} = \ln|x-1| - \frac{1}{3}$$

$$y = \frac{1}{\frac{1}{3} - \ln(x-1)}$$

Note: This solution is valid for  $1 < x < 1 + e^{1/3}$ .

2 :  $\left\{ \begin{array}{l} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \end{array} \right.$

2 :  $\left\{ \begin{array}{l} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{array} \right.$


5 :  $\left\{ \begin{array}{l} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration and} \\ \quad \text{uses initial condition} \\ 1 : \text{solves for } y \end{array} \right.$

Note: max 3/5 [1-2-0-0] if no constant of integration

Note: 0/5 if no separation of variables

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC**  
**2016 SCORING GUIDELINES**

**Question 5**



The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height  $h$ , the radius of the funnel is given by  $r = \frac{1}{20}(3 + h^2)$ , where  $0 \leq h \leq 10$ . The units of  $r$  and  $h$  are inches.

- (a) Find the average value of the radius of the funnel.
- (b) Find the volume of the funnel.
- (c) The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is  $h = 3$  inches, the radius of the surface of the liquid is decreasing at a rate of  $\frac{1}{5}$  inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?

$$\begin{aligned} \text{(a) Average radius} &= \frac{1}{10} \int_0^{10} \frac{1}{20}(3 + h^2) dh = \frac{1}{200} \left[ 3h + \frac{h^3}{3} \right]_0^{10} \\ &= \frac{1}{200} \left( \left( 30 + \frac{1000}{3} \right) - 0 \right) = \frac{109}{60} \text{ in} \end{aligned}$$

3 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^{10} \left( \frac{1}{20}(3 + h^2) \right)^2 dh = \frac{\pi}{400} \int_0^{10} (9 + 6h^2 + h^4) dh \\ &= \frac{\pi}{400} \left[ 9h + 2h^3 + \frac{h^5}{5} \right]_0^{10} \\ &= \frac{\pi}{400} \left( \left( 90 + 2000 + \frac{100000}{5} \right) - 0 \right) = \frac{2209\pi}{40} \text{ in}^3 \end{aligned}$$

3 :  $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(c) } \frac{dr}{dt} &= \frac{1}{20}(2h) \frac{dh}{dt} \\ \frac{1}{5} &= \frac{3}{10} \frac{dh}{dt} \\ \frac{dh}{dt} &= \frac{1}{5} \cdot \frac{10}{3} = \frac{2}{3} \text{ in/sec} \end{aligned}$$


3 :  $\begin{cases} 2 : \text{chain rule} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2015 SCORING GUIDELINES**

**Question 4**

Consider the differential equation  $\frac{dy}{dx} = 2x - y$ .

- (a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.
- (b) Find  $\frac{d^2y}{dx^2}$  in terms of  $x$  and  $y$ . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let  $y = f(x)$  be the particular solution to the differential equation with the initial condition  $f(2) = 3$ . Does  $f$  have a relative minimum, a relative maximum, or neither at  $x = 2$ ? Justify your answer.
- (d) Find the values of the constants  $m$  and  $b$  for which  $y = mx + b$  is a solution to the differential equation.



2 :  $\left\{ \begin{array}{l} 1 : \text{slopes where } x = 0 \\ 1 : \text{slopes where } x = 1 \end{array} \right.$

(b)  $\frac{d^2y}{dx^2} = 2 - \frac{dy}{dx} = 2 - (2x - y) = 2 - 2x + y$

In Quadrant II,  $x < 0$  and  $y > 0$ , so  $2 - 2x + y > 0$ .  
Therefore, all solution curves are concave up in Quadrant II.

2 :  $\left\{ \begin{array}{l} 1 : \frac{d^2y}{dx^2} \\ 1 : \text{concave up with reason} \end{array} \right.$

(c)  $\left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} = 2(2) - 3 = 1 \neq 0$

Therefore,  $f$  has neither a relative minimum nor a relative maximum at  $x = 2$ .

2 :  $\left\{ \begin{array}{l} 1 : \text{considers } \left. \frac{dy}{dx} \right|_{(x,y)=(2,3)} \\ 1 : \text{conclusion with justification} \end{array} \right.$

(d)  $y = mx + b \Rightarrow \frac{dy}{dx} = \frac{d}{dx}(mx + b) = m$

$$2x - y = m$$

$$2x - (mx + b) = m$$

$$(2 - m)x - (m + b) = 0$$

$$2 - m = 0 \Rightarrow m = 2$$

$$b = -m \Rightarrow b = -2$$

Therefore,  $m = 2$  and  $b = -2$ .

3 :  $\left\{ \begin{array}{l} 1 : \frac{d}{dx}(mx + b) = m \\ 1 : 2x - y = m \\ 1 : \text{answer} \end{array} \right.$


**AP<sup>®</sup> CALCULUS AB  
2014 SCORING GUIDELINES**

**Question 6**

Consider the differential equation  $\frac{dy}{dx} = (3 - y)\cos x$ . Let  $y = f(x)$  be the particular solution to the differential equation with the initial condition  $f(0) = 1$ . The function  $f$  is defined for all real numbers.

- (a) A portion of the slope field of the differential equation is given below. Sketch the solution curve through the point  $(0, 1)$ .
- (b) Write an equation for the line tangent to the solution curve in part (a) at the point  $(0, 1)$ . Use the equation to approximate  $f(0.2)$ .
- (c) Find  $y = f(x)$ , the particular solution to the differential equation with the initial condition  $f(0) = 1$ .

(a)



1 : solution curve

(b)  $\left. \frac{dy}{dx} \right|_{(x,y)=(0,1)} = 2 \cos 0 = 2$

An equation for the tangent line is  $y = 2x + 1$ .

$f(0.2) \approx 2(0.2) + 1 = 1.4$

2 :  $\left\{ \begin{array}{l} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{array} \right.$

(c)  $\frac{dy}{dx} = (3 - y)\cos x$

$\int \frac{dy}{3 - y} = \int \cos x \, dx$

$-\ln|3 - y| = \sin x + C$

$-\ln 2 = \sin 0 + C \Rightarrow C = -\ln 2$

$-\ln|3 - y| = \sin x - \ln 2$

Because  $y(0) = 1$ ,  $y < 3$ , so  $|3 - y| = 3 - y$

$3 - y = 2e^{-\sin x}$

$y = 3 - 2e^{-\sin x}$

Note: this solution is valid for all real numbers.

6 :  $\left\{ \begin{array}{l} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{array} \right.$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables

**AP<sup>®</sup> CALCULUS AB**  
**2013 SCORING GUIDELINES**

**Question 6**

Consider the differential equation  $\frac{dy}{dx} = e^y(3x^2 - 6x)$ . Let  $y = f(x)$  be the particular solution to the differential equation that passes through  $(1, 0)$ .

- (a) Write an equation for the line tangent to the graph of  $f$  at the point  $(1, 0)$ . Use the tangent line to approximate  $f(1.2)$ .
- (b) Find  $y = f(x)$ , the particular solution to the differential equation that passes through  $(1, 0)$ .

(a)  $\left. \frac{dy}{dx} \right|_{(x,y)=(1,0)} = e^0(3 \cdot 1^2 - 6 \cdot 1) = -3$

An equation for the tangent line is  $y = -3(x - 1)$ .

$f(1.2) \approx -3(1.2 - 1) = -0.6$

(b)  $\frac{dy}{e^y} = (3x^2 - 6x) dx$

$\int \frac{dy}{e^y} = \int (3x^2 - 6x) dx$

$-e^{-y} = x^3 - 3x^2 + C$

$-e^{-0} = 1^3 - 3 \cdot 1^2 + C \Rightarrow C = 1$

$-e^{-y} = x^3 - 3x^2 + 1$

$e^{-y} = -x^3 + 3x^2 - 1$

$-y = \ln(-x^3 + 3x^2 - 1)$

$y = -\ln(-x^3 + 3x^2 - 1)$

Note: This solution is valid on an interval containing  $x = 1$  for which  $-x^3 + 3x^2 - 1 > 0$ .

3 :  $\left\{ \begin{array}{l} 1 : \frac{dy}{dx} \text{ at the point } (x, y) = (1, 0) \\ 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{array} \right.$

6 :  $\left\{ \begin{array}{l} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{array} \right.$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables

**AP<sup>®</sup> CALCULUS AB**  
**2012 SCORING GUIDELINES**


**Question 5**

The rate at which a baby bird gains weight is proportional to the difference between its adult weight and its current weight. At time  $t = 0$ , when the bird is first weighed, its weight is 20 grams. If  $B(t)$  is the weight of the bird, in grams, at time  $t$  days after it is first weighed, then

$$\frac{dB}{dt} = \frac{1}{5}(100 - B).$$

Let  $y = B(t)$  be the solution to the differential equation above with initial condition  $B(0) = 20$ .

- (a) Is the bird gaining weight faster when it weighs 40 grams or when it weighs 70 grams? Explain your reasoning.
- (b) Find  $\frac{d^2B}{dt^2}$  in terms of  $B$ . Use  $\frac{d^2B}{dt^2}$  to explain why the graph of  $B$  cannot resemble the following graph.
- (c) Use separation of variables to find  $y = B(t)$ , the particular solution to the differential equation with initial condition  $B(0) = 20$ .



(a)  $\left. \frac{dB}{dt} \right|_{B=40} = \frac{1}{5}(60) = 12$

$$\left. \frac{dB}{dt} \right|_{B=70} = \frac{1}{5}(30) = 6$$

Because  $\left. \frac{dB}{dt} \right|_{B=40} > \left. \frac{dB}{dt} \right|_{B=70}$ , the bird is gaining weight faster when it weighs 40 grams.

(b)  $\frac{d^2B}{dt^2} = -\frac{1}{5} \frac{dB}{dt} = -\frac{1}{5} \cdot \frac{1}{5}(100 - B) = -\frac{1}{25}(100 - B)$

Therefore, the graph of  $B$  is concave down for  $20 \leq B < 100$ . A portion of the given graph is concave up.

(c)  $\frac{dB}{dt} = \frac{1}{5}(100 - B)$

$$\int \frac{1}{100 - B} dB = \int \frac{1}{5} dt$$

$$-\ln|100 - B| = \frac{1}{5}t + C$$

Because  $20 \leq B < 100$ ,  $|100 - B| = 100 - B$ .

$$-\ln(100 - 20) = \frac{1}{5}(0) + C \Rightarrow -\ln(80) = C$$

$$100 - B = 80e^{-t/5}$$

$$B(t) = 100 - 80e^{-t/5}, \quad t \geq 0$$

2 :  $\left\{ \begin{array}{l} 1 : \text{uses } \frac{dB}{dt} \\ 1 : \text{answer with reason} \end{array} \right.$

2 :  $\left\{ \begin{array}{l} 1 : \frac{d^2B}{dt^2} \text{ in terms of } B \\ 1 : \text{explanation} \end{array} \right.$

5 :  $\left\{ \begin{array}{l} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } B \end{array} \right.$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables



**AP<sup>®</sup> CALCULUS AB**  
**2011 SCORING GUIDELINES**

**Question 5**

At the beginning of 2010, a landfill contained 1400 tons of solid waste. The increasing function  $W$  models the total amount of solid waste stored at the landfill. Planners estimate that  $W$  will satisfy the differential equation  $\frac{dW}{dt} = \frac{1}{25}(W - 300)$  for the next 20 years.  $W$  is measured in tons, and  $t$  is measured in years from the start of 2010.

- (a) Use the line tangent to the graph of  $W$  at  $t = 0$  to approximate the amount of solid waste that the landfill contains at the end of the first 3 months of 2010 (time  $t = \frac{1}{4}$ ).
- (b) Find  $\frac{d^2W}{dt^2}$  in terms of  $W$ . Use  $\frac{d^2W}{dt^2}$  to determine whether your answer in part (a) is an underestimate or an overestimate of the amount of solid waste that the landfill contains at time  $t = \frac{1}{4}$ .
- (c) Find the particular solution  $W = W(t)$  to the differential equation  $\frac{dW}{dt} = \frac{1}{25}(W - 300)$  with initial condition  $W(0) = 1400$ .

(a)  $\left. \frac{dW}{dt} \right|_{t=0} = \frac{1}{25}(W(0) - 300) = \frac{1}{25}(1400 - 300) = 44$

The tangent line is  $y = 1400 + 44t$ .

$$W\left(\frac{1}{4}\right) \approx 1400 + 44\left(\frac{1}{4}\right) = 1411 \text{ tons}$$

$$2 : \begin{cases} 1 : \frac{dW}{dt} \text{ at } t = 0 \\ 1 : \text{answer} \end{cases}$$

(b)  $\frac{d^2W}{dt^2} = \frac{1}{25} \frac{dW}{dt} = \frac{1}{625}(W - 300)$  and  $W \geq 1400$

Therefore  $\frac{d^2W}{dt^2} > 0$  on the interval  $0 \leq t \leq \frac{1}{4}$ .

The answer in part (a) is an underestimate.

$$2 : \begin{cases} 1 : \frac{d^2W}{dt^2} \\ 1 : \text{answer with reason} \end{cases}$$

(c)  $\frac{dW}{dt} = \frac{1}{25}(W - 300)$

$$\int \frac{1}{W - 300} dW = \int \frac{1}{25} dt$$

$$\ln|W - 300| = \frac{1}{25}t + C$$

$$\ln(1400 - 300) = \frac{1}{25}(0) + C \Rightarrow \ln(1100) = C$$

$$W - 300 = 1100e^{\frac{1}{25}t}$$

$$W(t) = 300 + 1100e^{\frac{1}{25}t}, \quad 0 \leq t \leq 20$$

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } W \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

**AP<sup>®</sup> CALCULUS AB**  
**2008 SCORING GUIDELINES**

**Question 3**

Oil is leaking from a pipeline on the surface of a lake and forms an oil slick whose volume increases at a constant rate of 2000 cubic centimeters per minute. The oil slick takes the form of a right circular cylinder with both its radius and height changing with time. (Note: The volume  $V$  of a right circular cylinder with radius  $r$  and height  $h$  is given by  $V = \pi r^2 h$ .)

- (a) At the instant when the radius of the oil slick is 100 centimeters and the height is 0.5 centimeter, the radius is increasing at the rate of 2.5 centimeters per minute. At this instant, what is the rate of change of the height of the oil slick with respect to time, in centimeters per minute?
- (b) A recovery device arrives on the scene and begins removing oil. The rate at which oil is removed is  $R(t) = 400\sqrt{t}$  cubic centimeters per minute, where  $t$  is the time in minutes since the device began working. Oil continues to leak at the rate of 2000 cubic centimeters per minute. Find the time  $t$  when the oil slick reaches its maximum volume. Justify your answer.
- (c) By the time the recovery device began removing oil, 60,000 cubic centimeters of oil had already leaked. Write, but do not evaluate, an expression involving an integral that gives the volume of oil at the time found in part (b).

- (a) When  $r = 100$  cm and  $h = 0.5$  cm,  $\frac{dV}{dt} = 2000$  cm<sup>3</sup>/min  
and  $\frac{dr}{dt} = 2.5$  cm/min.

$$\begin{aligned}\frac{dV}{dt} &= 2\pi r \frac{dr}{dt} h + \pi r^2 \frac{dh}{dt} \\ 2000 &= 2\pi(100)(2.5)(0.5) + \pi(100)^2 \frac{dh}{dt} \\ \frac{dh}{dt} &= 0.038 \text{ or } 0.039 \text{ cm/min}\end{aligned}$$

$$4 : \begin{cases} 1 : \frac{dV}{dt} = 2000 \text{ and } \frac{dr}{dt} = 2.5 \\ 2 : \text{expression for } \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$$

- (b)  $\frac{dV}{dt} = 2000 - R(t)$ , so  $\frac{dV}{dt} = 0$  when  $R(t) = 2000$ .  
This occurs when  $t = 25$  minutes.  
Since  $\frac{dV}{dt} > 0$  for  $0 < t < 25$  and  $\frac{dV}{dt} < 0$  for  $t > 25$ ,  
the oil slick reaches its maximum volume 25 minutes after the device begins working.

$$3 : \begin{cases} 1 : R(t) = 2000 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

- (c) The volume of oil, in cm<sup>3</sup>, in the slick at time  $t = 25$  minutes is given by  $60,000 + \int_0^{25} (2000 - R(t)) dt$ .



$$2 : \begin{cases} 1 : \text{limits and initial condition} \\ 1 : \text{integrand} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB**  
**2008 SCORING GUIDELINES**

**Question 5**

Consider the differential equation  $\frac{dy}{dx} = \frac{y-1}{x^2}$ , where  $x \neq 0$ .

- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.  
(Note: Use the axes provided in the exam booklet.)
- (b) Find the particular solution  $y = f(x)$  to the differential equation with the initial condition  $f(2) = 0$ .
- (c) For the particular solution  $y = f(x)$  described in part (b), find  $\lim_{x \rightarrow \infty} f(x)$ .



2 :  $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{all other slopes} \end{cases}$

(b)  $\frac{1}{y-1} dy = \frac{1}{x^2} dx$   
 $\ln|y-1| = -\frac{1}{x} + C$   
 $|y-1| = e^{-\frac{1}{x} + C}$   
 $|y-1| = e^C e^{-\frac{1}{x}}$   
 $y-1 = ke^{-\frac{1}{x}}$ , where  $k = \pm e^C$   
 $-1 = ke^{-\frac{1}{2}}$   
 $k = -e^{\frac{1}{2}}$   
 $f(x) = 1 - e^{\left(\frac{1}{2} - \frac{1}{x}\right)}$ ,  $x > 0$

6 :  $\begin{cases} 1 : \text{separates variables} \\ 2 : \text{antidifferentiates} \\ 1 : \text{includes constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables

(c)  $\lim_{x \rightarrow \infty} 1 - e^{\left(\frac{1}{2} - \frac{1}{x}\right)} = 1 - \sqrt{e}$

1 : limit

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 5**

|                              |     |     |     |     |     |     |
|------------------------------|-----|-----|-----|-----|-----|-----|
| $t$<br>(minutes)             | 0   | 2   | 5   | 7   | 11  | 12  |
| $r'(t)$<br>(feet per minute) | 5.7 | 4.0 | 2.0 | 1.2 | 0.6 | 0.5 |

The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function  $r$  of time  $t$ , where  $t$  is measured in minutes. For  $0 < t < 12$ , the graph of  $r$  is concave down. The table above gives selected values of the rate of change,  $r'(t)$ , of the radius of the balloon over the time interval  $0 \leq t \leq 12$ . The radius of the balloon is 30 feet when  $t = 5$ . (Note: The volume of a sphere of radius  $r$  is given by  $V = \frac{4}{3}\pi r^3$ .)

- (a) Estimate the radius of the balloon when  $t = 5.4$  using the tangent line approximation at  $t = 5$ . Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when  $t = 5$ . Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate  $\int_0^{12} r'(t) dt$ . Using correct units, explain the meaning of  $\int_0^{12} r'(t) dt$  in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than  $\int_0^{12} r'(t) dt$ ? Give a reason for your answer.

(a)  $r(5.4) = r(5) + r'(5)\Delta t = 30 + 2(0.4) = 30.8$  ft  
Since the graph of  $r$  is concave down on the interval  $5 < t < 5.4$ , this estimate is greater than  $r(5.4)$ .

(b)  $\frac{dV}{dt} = 3\left(\frac{4}{3}\right)\pi r^2 \frac{dr}{dt}$   
 $\left.\frac{dV}{dt}\right|_{t=5} = 4\pi(30)^2(2) = 7200\pi$  ft<sup>3</sup>/min

(c)  $\int_0^{12} r'(t) dt \approx 2(4.0) + 3(2.0) + 2(1.2) + 4(0.6) + 1(0.5)$   
 $= 19.3$  ft  
 $\int_0^{12} r'(t) dt$  is the change in the radius, in feet, from  $t = 0$  to  $t = 12$  minutes.

(d) Since  $r$  is concave down,  $r'$  is decreasing on  $0 < t < 12$ . Therefore, this approximation, 19.3 ft, is less than  $\int_0^{12} r'(t) dt$ .

Units of ft<sup>3</sup>/min in part (b) and ft in part (c)

2 :  $\begin{cases} 1 : \text{estimate} \\ 1 : \text{conclusion with reason} \end{cases}$

3 :  $\begin{cases} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{cases}$

2 :  $\begin{cases} 1 : \text{approximation} \\ 1 : \text{explanation} \end{cases}$

1 : conclusion with reason

1 : units in (b) and (c)




**AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES**

**Question 5**


Consider the differential equation  $\frac{dy}{dx} = \frac{1+y}{x}$ , where  $x \neq 0$ .

- (a) On the axes provided, sketch a slope field for the given differential equation at the eight points indicated.  
(Note: Use the axes provided in the pink exam booklet.)



- (b) Find the particular solution  $y = f(x)$  to the differential equation with the initial condition  $f(-1) = 1$  and state its domain.

(a)



2 : sign of slope at each point and relative steepness of slope lines in rows and columns

(b)  $\frac{1}{1+y} dy = \frac{1}{x} dx$

$$\ln|1+y| = \ln|x| + K$$

$$|1+y| = e^{\ln|x| + K}$$

$$1+y = C|x|$$

$$2 = C$$

$$1+y = 2|x|$$

$$y = 2|x| - 1 \text{ and } x < 0$$

or

$$y = -2x - 1 \text{ and } x < 0$$

6 :  $\left\{ \begin{array}{l} 1 : \text{separates variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{array} \right.$

7 :  $\left\{ \begin{array}{l} \text{Note: max } 3/6 [1-2-0-0-0] \text{ if no} \\ \text{constant of integration} \\ \text{Note: } 0/6 \text{ if no separation of variables} \\ 1 : \text{domain} \end{array} \right.$


**AP Calculus AB**  
**Grading Rubric**  
**Area and Volume**

**AP<sup>®</sup> CALCULUS AB**  
**2015 SCORING GUIDELINES**

**Question 2**

Let  $f$  and  $g$  be the functions defined by  $f(x) = 1 + x + e^{x^2 - 2x}$  and  $g(x) = x^4 - 6.5x^2 + 6x + 2$ . Let  $R$  and  $S$  be the two regions enclosed by the graphs of  $f$  and  $g$  shown in the figure above.

- (a) Find the sum of the areas of regions  $R$  and  $S$ .
- (b) Region  $S$  is the base of a solid whose cross sections perpendicular to the  $x$ -axis are squares. Find the volume of the solid.
- (c) Let  $h$  be the vertical distance between the graphs of  $f$  and  $g$  in region  $S$ . Find the rate at which  $h$  changes with respect to  $x$  when  $x = 1.8$ .



- (a) The graphs of  $y = f(x)$  and  $y = g(x)$  intersect in the first quadrant at the points  $(0, 2)$ ,  $(2, 4)$ , and  $(A, B) = (1.032832, 2.401108)$ .

$$\begin{aligned} \text{Area} &= \int_0^A [g(x) - f(x)] dx + \int_A^2 [f(x) - g(x)] dx \\ &= 0.997427 + 1.006919 = 2.004 \end{aligned}$$

- (b) Volume =  $\int_A^2 [f(x) - g(x)]^2 dx = 1.283$

- (c)  $h(x) = f(x) - g(x)$   
 $h'(x) = f'(x) - g'(x)$   
 $h'(1.8) = f'(1.8) - g'(1.8) = -3.812$  (or  $-3.811$ )

4 :  $\left\{ \begin{array}{l} 1 : \text{limits} \\ 2 : \text{integrands} \\ 1 : \text{answer} \end{array} \right.$

3 :  $\left\{ \begin{array}{l} 2 : \text{integrand} \\ 1 : \text{answer} \end{array} \right.$

2 :  $\left\{ \begin{array}{l} 1 : \text{considers } h' \\ 1 : \text{answer} \end{array} \right.$




**AP<sup>®</sup> CALCULUS AB  
2014 SCORING GUIDELINES**

**Question 2**

Let  $R$  be the region enclosed by the graph of  $f(x) = x^4 - 2.3x^3 + 4$  and the horizontal line  $y = 4$ , as shown in the figure above.

- (a) Find the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = -2$ .
- (b) Region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is an isosceles right triangle with a leg in  $R$ . Find the volume of the solid.
- (c) The vertical line  $x = k$  divides  $R$  into two regions with equal areas. Write, but do not solve, an equation involving integral expressions whose solution gives the value  $k$ .



(a)  $f(x) = 4 \Rightarrow x = 0, 2.3$

$$\begin{aligned} \text{Volume} &= \pi \int_0^{2.3} [(4 + 2)^2 - (f(x) + 2)^2] dx \\ &= 98.868 \text{ (or } 98.867) \end{aligned}$$

4 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{answer} \end{cases}$

(b)  $\text{Volume} = \int_0^{2.3} \frac{1}{2} (4 - f(x))^2 dx$   
 $= 3.574 \text{ (or } 3.573)$

3 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$


(c)  $\int_0^k (4 - f(x)) dx = \int_k^{2.3} (4 - f(x)) dx$

2 :  $\begin{cases} 1 : \text{area of one region} \\ 1 : \text{equation} \end{cases}$

**AP<sup>®</sup> CALCULUS AB  
2013 SCORING GUIDELINES**

**Question 5**

Let  $f(x) = 2x^2 - 6x + 4$  and  $g(x) = 4\cos\left(\frac{1}{4}\pi x\right)$ . Let  $R$  be the region bounded by the graphs of  $f$  and  $g$ , as shown in the figure above.



- (a) Find the area of  $R$ .
- (b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = 4$ .
- (c) The region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is a square. Write, but do not evaluate, an integral expression that gives the volume of the solid.

$$\begin{aligned} \text{(a) Area} &= \int_0^2 [g(x) - f(x)] dx \\ &= \int_0^2 \left[ 4\cos\left(\frac{\pi}{4}x\right) - (2x^2 - 6x + 4) \right] dx \\ &= \left[ 4 \cdot \frac{4}{\pi} \sin\left(\frac{\pi}{4}x\right) - \left( \frac{2x^3}{3} - 3x^2 + 4x \right) \right]_0^2 \\ &= \frac{16}{\pi} - \left( \frac{16}{3} - 12 + 8 \right) = \frac{16}{\pi} - \frac{4}{3} \end{aligned}$$

4 :  $\begin{cases} 1 : \text{integrand} \\ 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b) Volume} &= \pi \int_0^2 \left[ (4 - f(x))^2 - (4 - g(x))^2 \right] dx \\ &= \pi \int_0^2 \left[ \left( 4 - (2x^2 - 6x + 4) \right)^2 - \left( 4 - 4\cos\left(\frac{\pi}{4}x\right) \right)^2 \right] dx \end{aligned}$$

3 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits and constant} \end{cases}$


$$\begin{aligned} \text{(c) Volume} &= \int_0^2 [g(x) - f(x)]^2 dx \\ &= \int_0^2 \left[ 4\cos\left(\frac{\pi}{4}x\right) - (2x^2 - 6x + 4) \right]^2 dx \end{aligned}$$

2 :  $\begin{cases} 1 : \text{integrand} \\ 1 : \text{limits and constant} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2012 SCORING GUIDELINES**

**Question 2**

Let  $R$  be the region in the first quadrant bounded by the  $x$ -axis and the graphs of  $y = \ln x$  and  $y = 5 - x$ , as shown in the figure above.



- (a) Find the area of  $R$ .
- (b) Region  $R$  is the base of a solid. For the solid, each cross section perpendicular to the  $x$ -axis is a square. Write, but do not evaluate, an expression involving one or more integrals that gives the volume of the solid.
- (c) The horizontal line  $y = k$  divides  $R$  into two regions of equal area. Write, but do not solve, an equation involving one or more integrals whose solution gives the value of  $k$ .

$$\ln x = 5 - x \Rightarrow x = 3.69344$$

Therefore, the graphs of  $y = \ln x$  and  $y = 5 - x$  intersect in the first quadrant at the point  $(A, B) = (3.69344, 1.30656)$ .

$$\begin{aligned} \text{(a) Area} &= \int_0^B (5 - y - e^y) dy \\ &= 2.986 \text{ (or } 2.985) \end{aligned}$$

OR

$$\begin{aligned} \text{Area} &= \int_1^A \ln x \, dx + \int_A^5 (5 - x) \, dx \\ &= 2.986 \text{ (or } 2.985) \end{aligned}$$

$$\text{(b) Volume} = \int_1^A (\ln x)^2 \, dx + \int_A^5 (5 - x)^2 \, dx$$

$$\text{(c) } \int_0^k (5 - y - e^y) \, dy = \frac{1}{2} \cdot 2.986 \text{ (or } \frac{1}{2} \cdot 2.985)$$

3 :  $\left\{ \begin{array}{l} 1 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{answer} \end{array} \right.$

3 :  $\left\{ \begin{array}{l} 2 : \text{integrands} \\ 1 : \text{expression for total volume} \end{array} \right.$


3 :  $\left\{ \begin{array}{l} 1 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{equation} \end{array} \right.$

**AP<sup>®</sup> CALCULUS AB**  
**2011 SCORING GUIDELINES**

**Question 3**

Let  $R$  be the region in the first quadrant enclosed by the graphs of  $f(x) = 8x^3$  and  $g(x) = \sin(\pi x)$ , as shown in the figure above.

- (a) Write an equation for the line tangent to the graph of  $f$  at  $x = \frac{1}{2}$ .
- (b) Find the area of  $R$ .
- (c) Write, but do not evaluate, an integral expression for the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = 1$ .



(a)  $f\left(\frac{1}{2}\right) = 1$   
 $f'(x) = 24x^2$ , so  $f'\left(\frac{1}{2}\right) = 6$

An equation for the tangent line is  $y = 1 + 6\left(x - \frac{1}{2}\right)$ .

(b) Area =  $\int_0^{1/2} (g(x) - f(x)) dx$   
 $= \int_0^{1/2} (\sin(\pi x) - 8x^3) dx$   
 $= \left[ -\frac{1}{\pi} \cos(\pi x) - 2x^4 \right]_{x=0}^{x=1/2}$   
 $= -\frac{1}{8} + \frac{1}{\pi}$

(c)  $\pi \int_0^{1/2} \left( (1 - f(x))^2 - (1 - g(x))^2 \right) dx$   
 $= \pi \int_0^{1/2} \left( (1 - 8x^3)^2 - (1 - \sin(\pi x))^2 \right) dx$


2 :  $\begin{cases} 1 : f'\left(\frac{1}{2}\right) \\ 1 : \text{answer} \end{cases}$

4 :  $\begin{cases} 1 : \text{integrand} \\ 2 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

3 :  $\begin{cases} 1 : \text{limits and constant} \\ 2 : \text{integrand} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2010 SCORING GUIDELINES**

**Question 4**



Let  $R$  be the region in the first quadrant bounded by the graph of  $y = 2\sqrt{x}$ , the horizontal line  $y = 6$ , and the  $y$ -axis, as shown in the figure above.

- (a) Find the area of  $R$ .
- (b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = 7$ .
- (c) Region  $R$  is the base of a solid. For each  $y$ , where  $0 \leq y \leq 6$ , the cross section of the solid taken perpendicular to the  $y$ -axis is a rectangle whose height is 3 times the length of its base in region  $R$ . Write, but do not evaluate, an integral expression that gives the volume of the solid.

(a) 
$$\text{Area} = \int_0^9 (6 - 2\sqrt{x}) \, dx = \left( 6x - \frac{4}{3}x^{3/2} \right) \Big|_{x=0}^{x=9} = 18$$

3 :  $\begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

(b) 
$$\text{Volume} = \pi \int_0^9 \left( (7 - 2\sqrt{x})^2 - (7 - 6)^2 \right) dx$$

3 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits and constant} \end{cases}$

(c) Solving  $y = 2\sqrt{x}$  for  $x$  yields  $x = \frac{y^2}{4}$ .

Each rectangular cross section has area  $\left( 3 \frac{y^2}{4} \right) \left( \frac{y^2}{4} \right) = \frac{3}{16}y^4$ .


$$\text{Volume} = \int_0^6 \frac{3}{16}y^4 \, dy$$

3 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB**  
**2009 SCORING GUIDELINES**

**Question 4**

Let  $R$  be the region in the first quadrant enclosed by the graphs of  $y = 2x$  and  $y = x^2$ , as shown in the figure above.



- (a) Find the area of  $R$ .
- (b) The region  $R$  is the base of a solid. For this solid, at each  $x$  the cross section perpendicular to the  $x$ -axis has area  $A(x) = \sin\left(\frac{\pi}{2}x\right)$ . Find the volume of the solid.
- (c) Another solid has the same base  $R$ . For this solid, the cross sections perpendicular to the  $y$ -axis are squares. Write, but do not evaluate, an integral expression for the volume of the solid.

$$\begin{aligned} \text{(a) Area} &= \int_0^2 (2x - x^2) dx \\ &= x^2 - \frac{1}{3}x^3 \Big|_{x=0}^{x=2} \\ &= \frac{4}{3} \end{aligned}$$

$$3 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$\begin{aligned} \text{(b) Volume} &= \int_0^2 \sin\left(\frac{\pi}{2}x\right) dx \\ &= -\frac{2}{\pi} \cos\left(\frac{\pi}{2}x\right) \Big|_{x=0}^{x=2} \\ &= \frac{4}{\pi} \end{aligned}$$


$$3 : \begin{cases} 1 : \text{integrand} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

$$\text{(c) Volume} = \int_0^4 \left(\sqrt{y} - \frac{y}{2}\right)^2 dy$$

$$3 : \begin{cases} 2 : \text{integrand} \\ 1 : \text{limits} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB**  
**2008 SCORING GUIDELINES**

**Question 1**



Let  $R$  be the region bounded by the graphs of  $y = \sin(\pi x)$  and  $y = x^3 - 4x$ , as shown in the figure above.

- Find the area of  $R$ .
- The horizontal line  $y = -2$  splits the region  $R$  into two parts. Write, but do not evaluate, an integral expression for the area of the part of  $R$  that is below this horizontal line.
- The region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is a square. Find the volume of this solid.
- The region  $R$  models the surface of a small pond. At all points in  $R$  at a distance  $x$  from the  $y$ -axis, the depth of the water is given by  $h(x) = 3 - x$ . Find the volume of water in the pond.

(a)  $\sin(\pi x) = x^3 - 4x$  at  $x = 0$  and  $x = 2$

$$\text{Area} = \int_0^2 (\sin(\pi x) - (x^3 - 4x)) dx = 4$$

3 : { 1 : limits  
1 : integrand  
1 : answer

(b)  $x^3 - 4x = -2$  at  $r = 0.5391889$  and  $s = 1.6751309$

$$\text{The area of the stated region is } \int_r^s (-2 - (x^3 - 4x)) dx$$

2 : { 1 : limits  
1 : integrand

(c)  $\text{Volume} = \int_0^2 (\sin(\pi x) - (x^3 - 4x))^2 dx = 9.978$

2 : { 1 : integrand  
1 : answer


(d)  $\text{Volume} = \int_0^2 (3 - x)(\sin(\pi x) - (x^3 - 4x)) dx = 8.369$  or  $8.370$

2 : { 1 : integrand  
1 : answer

AP<sup>®</sup> CALCULUS AB  
2006 SCORING GUIDELINES (Form B)

Question 2

Let  $f$  be the function defined for  $x \geq 0$  with  $f(0) = 5$  and  $f'$ , the first derivative of  $f$ , given by  $f'(x) = e^{(-x/4)} \sin(x^2)$ . The graph of  $y = f'(x)$  is shown above.



- (a) Use the graph of  $f'$  to determine whether the graph of  $f$  is concave up, concave down, or neither on the interval  $1.7 < x < 1.9$ . Explain your reasoning.
- (b) On the interval  $0 \leq x \leq 3$ , find the value of  $x$  at which  $f$  has an absolute maximum. Justify your answer.
- (c) Write an equation for the line tangent to the graph of  $f$  at  $x = 2$ .

(a) On the interval  $1.7 < x < 1.9$ ,  $f'$  is decreasing and thus  $f$  is concave down on this interval.

2 : { 1 : answer  
1 : reason

(b)  $f'(x) = 0$  when  $x = 0, \sqrt{\pi}, \sqrt{2\pi}, \sqrt{3\pi}, \dots$   
On  $[0, 3]$   $f'$  changes from positive to negative only at  $\sqrt{\pi}$ . The absolute maximum must occur at  $x = \sqrt{\pi}$  or at an endpoint.

3 : { 1 : identifies  $\sqrt{\pi}$  and 3 as candidates  
- or -  
indicates that the graph of  $f$  increases, decreases, then increases  
1 : justifies  $f(\sqrt{\pi}) > f(3)$   
1 : answer

$$f(0) = 5$$

$$f(\sqrt{\pi}) = f(0) + \int_0^{\sqrt{\pi}} f'(x) dx = 5.67911$$

$$f(3) = f(0) + \int_0^3 f'(x) dx = 5.57893$$

This shows that  $f$  has an absolute maximum at  $x = \sqrt{\pi}$ .

(c)  $f(2) = f(0) + \int_0^2 f'(x) dx = 5.62342$

$$f'(2) = e^{-0.5} \sin(4) = -0.45902$$

$$y - 5.623 = (-0.459)(x - 2)$$

4 : { 2 :  $f(2)$  expression  
1 : integral  
1 : including  $f(0)$  term  
1 :  $f'(2)$   
1 : equation